

*Original Article*

**Different approaches of trans-cervical artificial insemination in oestrus  
synchronized indigenous field ewes using frozen ram semen**

Mohammad Asaduzzaman<sup>1,2</sup>, Pankaj Kumar Jha<sup>3</sup>, Amit Saha<sup>2</sup>, Md. Golam Shahi

Alam<sup>2</sup>, Farida Yeasmin Bari<sup>2\*</sup>

<sup>1</sup>Department of Livestock Services, Farmgate, Dhaka-1215, Bangladesh

<sup>2</sup>Department of Surgery and Obstetrics, Bangladesh Agricultural University,

Mymensingh-2204

<sup>3</sup>Animal Breeding Division, Nepal Agricultural Research Council, Lalitpur - 44703,

Nepal

**\* Corresponding author, Email address:** [faridabari06@gmail.com](mailto:faridabari06@gmail.com)

**Abstract**

The study aimed to determine the pregnancy rates of indigenous field sheep following different approaches to Trans-cervical insemination (TCAI). TCAI was performed in oestrus induced ewes using two straws of frozen-thawed semen. In group 1, single-time TCAI was performed on standing heat in traditional/ manual restrained ewes (n=20) and lab cradle restrained ewes (n=20). In group 2, TCAI was performed in manually restrained ewes single-time after 46-48h of PGF2a injection (n=20), and double-times after 46-48h & 52-56h of PGF2a injection (n=20). In group 3, TCAI was performed in manually restrained ewes single time after 52-56h of PGF2a injection and 20 minutes after oxytocin injection (n=10), and 20 minutes after oxytocin-estradiol benzoate injection (n=11). The Pregnancy rates were 35% and 45 % in traditional and cradle restrained TCAI, 35 and 55 % in single-time and double-times TCAI, and 50 and 55 % in oxytocin and oxytocin-estradiol cervix-treated TCAI, respectively. Although

the pregnancy rates were not significant ( $p>0.05$ ) within experimental groups, the TCAI after cervical treatment and double-time TCAI importantly had more pregnancy than other approaches of TCAI. Finally it can be concluded that double-times TCAI would be an effective and acceptable approach for field sheep artificial breeding using frozen semen.

**Keywords:** Sheep, Oestrus synchronization, Different approaches, Transcervical insemination

## 1. Introduction

Animals are bred artificially by incorporating semen into a natural or synchronized female's oestrus genital tract using instruments. Artificial insemination (AI), in no doubt, is the oldest technique within assisted reproduction (AR) (Falchi, Taema, La Clanche, & Scaramuzzi, 2012) used initially to avoid male injuries and prevent sexually transmitted diseases (Hernández et al., 2015). Soon it was recognized by farmers as a method of choice to add or spread the superior genetic value of males for enhancing the production of improved offspring with high productive traits and to control contagious diseases in animal herds (Evans & Maxell, 1987; Alvares, Cruz & Ferreira, 2015). It also facilitates maximum use of superior rams by lowering the sperm concentration in semen dose, increasing the number of insemination and quality offspring.

The animal oestrus cycle can be manipulated to bring them to heat within a predefined time frame of 36 to 96 hours by one or a combination of three hormones: prostaglandins, progesterone, and GnRH. It facilitates a group of animals to be synchronized quickly for efficient artificial insemination (AI) (Tsuma, Khan, Okeyo Mwai & Ibrahim, 2015). Islam (2011) stated that the cyclic female shows oestrus within

2 to 6 days following prostaglandin F2 $\alpha$  (Estrumate®, In-Synch®). [Gizaw et al., \(2016b\)](#) described that when a single injection of prostaglandin was given to a flock of cycling ewes, 60 to 70 percent of the flock will exhibit a synchronized estrous beginning 30 to 48 hours later ([Schoenian, 2012](#); [Tsuma, Khan, Okeyo Mwai & Ibrahim, 2015](#)). [Islam \(2011\)](#) stated that because not all stages of the estrous cycle were similarly receptive to treatment, a double injection system 7 days or 11 days apart, to make sure all functional CL regress was the most widely used approach in small ruminant.

The sheep AI is involved with appropriate knowledge and breeds ([Fonseca, Souza & Camargo, 2010](#)), female body condition, prolificacy, and oestrus status. In addition, farm conditions, time of year, season, the technique of AI, and technician skills also influence the fertility rate ([Anel et al., 2005](#)). Apart from these, the structure of the sheep cervix is another crucial barrier for transcervical AI (TCAI) with its 4-7 convoluted rings that prevent the passage of AI pipette through to the body of the uterus ([Evans & Maxell, 1987](#)). The role of the cervical canal in regulating frozen sperm transport in sheep is still problematic, despite various inseminating equipment that has been tried with cervix manipulation for effective insemination ([Aral, Temamogullari & Aral, 2011](#)). Even though artificial insemination with laparoscopy and laparotomy techniques has been shown to have compelling performances, high cost, time-consuming, and requirement of a particular type of equipment are the main drawbacks of their wider use ([Evans & Maxwell, 1987](#); [Torres & Sevellec, 1987](#)). Many to overcome all these factors, hormones, primarily, Oxytocin and prostaglandin E (PGE), were found to promote significant cervical dilatation ([Khalifa, Sayre & Lewis, 1992](#); [Wulster-Radcliffe, Costine & Lewis, 1999](#); [Masoudi et al., 2017](#)).

Azizunnesa, Zohara, Islam, Bari & Alam (2013), Azizunnesa, Bari & Alam (2014), and Zohara, Islam, Alam & Baric (2014) performed some baseline studies on sheep oestrus synchronization using different hormones protocol, ram semen preservation, and artificial insemination in the research station of Bangladesh Agricultural University, Mymensingh. Al Mansur et al., (2018) had little exposure to the performance of Transcervical in field indigenous ewes. The present study aimed to determine the pregnancy rates of indigenous field sheep following different approaches of TCAI.

## **2. Materials and Methods**

### **Study site, animal, and management**

The experiment was carried out on Bangladeshi indigenous Jamuna basin sheep of different field sheep farms at Gopalpur and Bhoapur Upazilla under Tangail district and Mymensingh Sadar Upazila under Mymensingh district during three subsequent breeding seasons from November 2018 to March 2021. All the ewes were mainly under the farmer's management systems, such as they were maintained on pasture from morning to dusk with free excess of safe drinking water. The sheep were 2–4 years aged and or at least one parity.

### **Sampling method and approval**

Sample sizes were calculated using a formula,  $n = N / 1 + N(e)^2$  (Yamane, 1967: 886), where  $n$  is the sample size,  $N$  is the population size, and  $e$  is the level of precision. From 20 field sheep farmers, a total of 200 non-pregnant indigenous ewes were randomly selected as the hypothesized population size for the current study. At 95 percent confidence level and  $P = 0.5$ , the calculated sample size was 134 ewes. All experimental protocols followed the Animal Welfare and Experimentation Ethics

Committee's (AWEEC) recommendations for animal care and used [No. AWEEC/BAU/2019 (54), dated: 26/12/2019].

## **Experimental design**

A total of 134 ewes were brought under the oestrus synchronization program, and oestrus was confirmed to 108 ewes. Transcervical artificial insemination was performed by dividing them into three experimental groups where 101 ewes were countable for efficient artificial insemination, and 7 ewes were excluded for defective artificial insemination.

### **Group 1: Techniques of TCAI (n=40)**

- Traditional restrained, and TCAI on standing heat (n = 20)
- Lab cradle restrained, and TCAI on standing heat (n = 20)

### **Group 2: Time of TCAI (n=40)**

- Single-time TCAI after 46-48h of PGF2a injection (n=20)
- Double time TCAI after 46-48h and 52-56h of PGF2a injection (n= 20)

### **Group 3: Cervix treatment and TCAI (n = 21)**

- TCAI after 52-56h of PGF2a and 20 minutes of Oxytocin (OT) injection (n=10)
- TCAI after 52-56h of PGF2a and 20 minutes of Oxytocin(OT)+ Oestradiol (OE) injection (n=11)

## **Semen preparation, thawing, and AI gun loading**

The selected semen samples were frozen according to a conventional laboratory standard protocol by Jha et al., (2019). After thawing the frozen straw, the post-thaw seminal parameters were measured (37°C for 30 seconds). While the production motility was at least 40%, they were stored. During TCAI, the straw was thawed as before; the company end was fitted to an AI gun (<sup>®</sup>Inseminator for small ruminants, 0.25 ml straws, Minitube). The straw's sealed end was cut-opened with clean scissors and finally loaded into a pipette for TCAI.

## Synchronization and detection of oestrus

Oestrus cycles of all groups were synchronized by prostaglandin (Gizaw et al., 2016) using two injection protocols of Rekha, Zohara, Bari & Alam (2016) at the unknown stage of the oestrus cycle of the non-pregnant ewes confirmed by ultrasonography (DRAMÍŃSKI ANIMAL *profi*, Poland). Two injections of Ovuprost® (Cloprostenol sterile injection, BOMAC Laboratories Ltd., New Zealand) were given at 9 days intervals i.m at the dose rate of 0.4 ml/ewe (Zohara, Alamc & Baric, 2014) to make sure all functional CL regress. The heat was proved to be in oestrus while an aproned ram mounted her within 24 to 48 h of prostaglandin injection.

Figure 1. (A) Oestrus detection by the aproned ram, (B) TCAI with traditional and manual restrained ewe, (C) TCAI with lab cradle restrained ewe,

## Traditional and lab cradle restrained TCAI approach

In a traditionally restrained approach, the farmer or assistant stood over the ewe holdings, securing her two hind legs, and elevated the hindquarter as the ewe was in a slanting position. With firm and gentle pressure, the lubricated (KLY® Jelly) vaginal speculum was carefully inserted into the full depth of the vagina (10-13cm) following the slope of the female's rump. A light source (pen torch) was attached with a speculum to assist the visual inspection and locate and position the cervical os by pulling back or rotating the speculum slightly. The straw-loaded AI gun was inserted and plunged the semen into the cervical os. Two straws of frozen semen were used in artificial insemination containing  $200 \times 10^6$  / 0.25 ml straw of frozen semen have at least 40% motile spermatozoa. While in restrained cradle approach, the ewe boarded to cradle bed keeping her hindquarters elevated with head down position with the body in dorsoventral position. The hind legs were fixed with an upper rail of the cradle using

ropes, and the front legs with two vertical rails. One assistant stood beside the cradle, holding and fixing the female head. The ewe was then inseminated following the traditional method.

#### **Single-time and double-times TCAI approach**

Single time transcervical artificial insemination approach was performed at 46 - 48 h of induction of oestrus, and the double-time approach was performed at 46 - 48 h and 52 - 56 h of induction of oestrus as it was close to Visser & Salamon's (1974) who performed insemination at 12-14 h, and 23-25 h of onset of oestrus. During TCAI, the ewes were restrained traditionally, as in the previous description of group 1.

#### **The cervix treatment approach of TCAI**

The synchronized ewe cervix was ripened with oxytocin (OT) (Linda-S® DS, Synthetic Oxytocin USP 10 IU/ml, Nuvista Pharma Limited, Gazipur, Bangladesh) at a dose of 50 IU i/m per animal 20 minutes before TCAI in one group of animals. In other groups, the synchronized ewe cervix was ripened by treating combine with Oxytocin (0.5ml, 2.5-10 IU/small ruminants) and oestradiol (1ml, 2mg/small ruminants) (Estradiol Benzoate Injection, QUANYU Sheng Wukeji, China) i/m per animal 20 minutes before insemination. The ewes were restrained as a manual or traditional approach before insemination.

#### **Pregnancy diagnosis**

The conception was detected by trans-abdominal ultrasonography after 45 to 60 days of insemination using an ultrasound scanner, DRAMIŃSKI ANIMAL *profi*, Poland.

#### **Data analysis**

Data are presented as percentages. The Chi-square statistic of SPSS var. 22 was used to assess differences among percentages of pregnancy rates in different methods of transcervical artificial insemination. The significance level was defined as a p-value < 0.05.

### 3. Results

The pregnancy rate of transcervical artificial insemination in three experiment groups of the animal following different approaches are presented in Table 1. The result of group 1 indicates the percentage of animals successfully impregnated or not concerning the restraining technique of animals during TCAI of indigenous ewes. The restraining techniques or methods had no impact on pregnancy rates ( $p>0.05$ ). Although the pregnancy rate was higher in cradle-restrained animals than in manual or traditional restrained animals, the difference was not statistically significant.

The result of group 2 represents the percentages of pregnancy rate of single-time and double-times TCAI in indigenous field sheep at (46 - 48) hand (46 - 48) & (54 - 56) h of oestrus induction. Double-times TCAI had a higher pregnancy rate than single-time TCAI. Although there was no significant ( $p>0.05$ ) difference, double-times TCAI in field sheep showed a highly effective pregnancy rate than single-time TCAI.

The result of group 3 shows the percentage of pregnancy rate of indigenous ewes following TCAI after treatment with cervical dilators: oxytocin (OT) and oxytocin and estradiol benzoate ( $E_2$ +OT) combined. The OT and  $E_2$ +OT treatment groups have a higher pregnancy rate (50 and 55%, respectively) than the traditional restrained TCAI approach group (35%). Although there were no statistical differences ( $p>0.05$ ) in pregnancy rates, the combined dilator ( $E_2$ +OT) showed more effectiveness than the single dilator (OT).



#### 4. Discussion

A successful technique or approach for successful artificial insemination has not yet been well established in sheep due to its cervical-specific anatomy compared to horses and cattle (Halbert, Dobson, Walton & Buckrell, 1990). The Guelph system for trans-cervical AI (GST-AI) has been developed; it is not suitable for farmer's application as it requires specially designed instruments and unique positioning of ewe and trained and experienced inseminator. Moreover, cervical retraction and penetration rate, injury, abscesses, infections, and poor pregnancy are the associated unappreciated factors of this technique (Buschbeck, 2020).

The current study performed TCAI in the farmer's field by traditionally or manually restraining the ewes. The lower pregnancy rate of 35% obtained from this approach was suggested by the observation of Gage (1994), Salamon & Maxwell (1995), and Donovan, Hanrahan, Kummen, Duffy & Boland (2004), as 33, 34.8, and 36%, respectively. To obtain the maximum benefit of TCAI, we performed trans-cervical artificial insemination by restraining and boarding the ewes in the lab cradle and observed a comparatively higher pregnancy rate (45%) than the manual restraining approach. The lower pregnancy rate in the manual restraining approach might be due to the difficulty of keeping the animal substantially immobile for a longer duration of AI. Sometimes, people fix the sheep less conveniently, and such instability condition even leads to injury, which lowers the success rate of artificial insemination. The higher pregnancy rate in the lab cradle restrained approach might be due to the stationary state of the animal, facilitating the operation insemination, and lifting the sheep bed and buttocks to prevent semen backflow, simultaneously improving the success rate of insemination of sheep (Google Patent, 2012). However, such a higher pregnancy rate

was in agreement with the observation of Maxwell & Hewitt (1986), Donovan et al. (2001), and Donovan, Hanrahan, Kummen, Duffy & Boland (2004) as of 42.2, 44, and 46%, respectively.

The time of AI is critical as it is related to ovulation time. Ovulation occurs at the end phase of oestrus. As such, AI needs to be done sufficient time before ovulation so that sperm can migrate to the fertilization site, the ampulla. In tradition, oestrus is observed twice (morning and evening) daily. While oestrus is observed in the morning, AI is performed in the evening (around 10 h of oestrus onset). Oestrus is observed in the evening; insemination is done in the morning (about 14 h of oestrus onset). The fresh semen has a fertilizing lifetime of more than 24 h, whether not more than 12 h for frozen semen in the female reproductive tract. Therefore, AI time needs to be adjusted according to semen types (Kumar & Naqvi, 2014).

Adjustment of AI time is also essential in the case of oestrus synchronization. Fernandez-Abella, Preve & Villegas (2003) suggested that the best time of AI is 46h, 48 to 72 h by Karagiannidis, Varsakeli, Karatzas & Brozos (2001), and 58 to 63 h after the pessary removal by Donovan et al., (2001), Donovan, Hanrahan, Kummen, Duffy & Boland (2004). Single-time TCAI of the present study was performed at (46 - 48) h of oestrus induction and double-times TCAI at (46 - 48) and (54 - 56) h of oestrus induction. The pregnancy rate of single-time TCAI (35%) was in agreement with the reported pregnancy rates of Gage (1994), Salamon & Maxwell (1995), and Donovan, Hanrahan, Kummen, Duffy & Boland (2004) as of 33%, 22.2 - 34.8%, and 36%, respectively. The higher conception rate of double-times TCAI (55%) was in line with the findings of Stefanov et al., (2006); Halbert, Dobson, Walton & Buckrell (1990), Wulster- Radcliffe, and Lewis (2002) as of 53.33%, 58%, and 59.3%, respectively.

Similarly, Kumar & Naqvi (2014) found a higher pregnancy rate of transcervical artificial insemination in double-times insemination than in single-time insemination (26.4 vs. 20%). Maxwell (1980) found a 29% conception rate after insemination at 50 and 60 h intervals. Halbert, Dobson, Walton & Buckrell (1990C); Buckrell et al., (1992) performed transcervical artificial insemination at 54 h intervals of synchronization and found conception rates as 24.4 - 50.7 and 34.7 - 56.7 %, respectively. Similarly, Smith, Parr, Beaumont & Oliver (1995) found 40% of the conception rate following trans-cervical artificial insemination at 50-56 h intervals of the synchronization.

The deeper intra cervix insemination in ewes facilitates more spermatozoa to reach the fertilization site and consequently increases the fertilization and pregnancy rates (Eppleston, Salamon, Moore & Evans 1994; Anel et al., 2006). However, sheep's cervical structure impedes these successful insemination functions (Wulster-Radcliffe, Costine & Lewis, 1999). Some of the procedures have included treatments to dilate the cervix and reduce the difficulty of manipulating AI catheters through the cervix and into the uterus (Wulster-Radcliffe, Wang, & Lewis, 2004; Anel et al., 2006; Candappa, Bainbridge, Price, Hourigan & Bartlewski, 2009). These are physical-mechanical methods based on physical stress to the cervix or transfer of sperm by a flexible or semi-flexible pipette (Wulster-Radcliffe & Lewis, 2002; Wulster-Radcliffe, Wang, & Lewis, 2004) that may injure the cervix and produce some anti-sperm secretions leading to lower sperm viability and consequently lower conception rate (Hawk, 1983; Sayre & Lewis, 1997). Even though laparoscopic and laparotomy methods of artificial inseminations are more effective, they come at a high cost and require much time and specialized equipment (Evans & Maxwell, 1987; Torres & Sevellec, 1987). As a result,

cervix treatment to improve sperm reach to the fertilization site by altering its physiological characteristics without causing injury is the most effective method of choice.

It has been shown that oxytocin and prostaglandin E (PGE) hormones effectively dilate the cervix and increase the pregnancy rate (Khalifa, Sayre, & Lewis, 1992; Wulster-Radcliffe, Costine & Lewis, 1999). In the luteal phase, they have no effect and result in minimal damage or injury when passages of AI pipettes into the cervix (Masoudi et al., 2017). Pipette passage through the cervix is facilitated by oxytocin, allowing for an intrauterine supply of semen. Oxytocin aids the pipette's passage past the cervix and sperm deposit within the uterus. It was also found that the processes of transcervical (TC) embryo transfer (ET) in sheep improved considerably, increasing cervical dilatation, exogenous estradiol-17 beta (E2), and oxytocin (OT) (Wulster-Radcliffe, Costine & Lewis, 1999). Masoudi et al., (2017) found the highest cervical dilation (90%) and (100%) in OT and E2+OT groups with the highest pregnancy of 60 and 65% of fat-tailed ewes, respectively. These findings were suggested with our present results of 50 and 55% for OT and E2+OT groups, respectively. But the level of variation between OT and E2+OT groups might be due to differences in sheep breeds, body sizes, the sources, and doses of dilators used in the present study. For the present study, we also observed more pregnancy rates while using combined dilators (OT+E2) than single OT dilators, and this might be due to more penetration effects as described by Masoudi et al., (2017) as E2+OT (4.10 cm) and OT (3.90 cm). While sheep cervix remains more dilated, deep cervical insemination produces higher fertilization. Fukui & Roberts (1978) also agreed with this report and observed a 45% fertility rate for deep cervical insemination with frozen semen (300

million sperm/ dose). In another report, Falchi, Taema, La Clanche & Scaramuzzi (2012) also stated that maximum cervical penetration in the preovulatory period with estradiol concentrations suggests the cervix's relaxation probably through an oxytocin-PGE mediated pathway.

## 5. Conclusions

With the application of frozen semen, the present research studied a more effective TCAI approach for artificial breeding in field ewes. Based on the findings, the following conclusions can be drawn:

- i. TCAI with the traditionally restrained approach is straightforward, less time-consuming, and more suitable to apply in field perspective than TCAI with a lab cradle restrained approach. Although a little higher pregnancy rate was obtained in the second stated approach, it requires a particular cradle instrument that is not suitable for carrying and instant application to field farms.
- ii. Oxytocin and estradiol treatment may permit routine transcervical artificial inseminations with a comparatively higher pregnancy rate. Still, the availability of these hormones, time-consuming effects, and smooth performance of insemination in many sheep may be significantly negative issues of adopting artificial insemination.
- iii. The time of TCAI was the most effective than other studied approaches. Double-times TCAI covered the ovulation and fertilization time and produced a higher conception rate than single-time TCAI and might be the approach of choice in field sheep artificial breeding.

## Acknowledgments

We are grateful to Upazilla Livestock Officer and Veterinary Field Assistant, Gopalpuras, Bhuapur, Tangail district, and reproduction lab assistant, BAU, Mymensingh.

## References

- Al Mansur, M. A., Alam, M. G. S., Jha, P. K., Rimon, M. A., Naher, N., & Bari, F. Y. (2018). Conception rate following intra-cervical artificial insemination using frozen semen at field level in indigenous sheep of Bangladesh. *Asian Journal of Medical and Biological Research*, 4(1), 55-62.
- Alvares, C. T. G., da Cruz, J. F., & Ferreira, M. L. (2015). Técnicas de inseminação artificial e implicações fisiopatológicas em ovinos. *PubVet*, 9, 195-251.
- Anel, L., Alvarez, M., Martinez- Pastor, F., Garcia- Macias, V., Anel, E., & De Paz, P. (2006). Improvement strategies in ovine artificial insemination. *Reproduction in Domestic Animals*, 41, 30-42.
- Anel, L., Kaabi, M., Abroug, B., Alvarez, M., Anel, E., Boixo, J. C., ... & De Paz, P. (2005). Factors influencing the success of vaginal and laparoscopic artificial insemination in churra ewes: a field assay. *Theriogenology*, 63(4), 1235-1247.
- Aral, F., Temamogullari, F., & Aral, S. S. (2011). Mechanical and pharmacologic applications of artificial insemination in ewes. *Veterinary Medicine and Science*, 243-254.
- Azizunnesa, A., Zohara, B. F., Bari, F. Y., & Alam, M. G. S. (2013). Effects of concentrate supplementation on reproductive performances and semen quality of indigenous rams in Bangladesh. *Journal of Embryo Transfer*, 28(4), 325-335.
- Azizunnesa, Z. B., Bari, F. Y., & Alam, M. G. S. (2014). Baseline study of reproductive performances of indigenous rams in Bangladesh. *Journal of Agriculture and Veterinary Science*, 7(6), 83-89.
- Buckrell, B. C., Buschbeck, C., Gartley, C. J., Kroetsch, T., McCutcheon, W., Martin, J., ... & Walton, J. S. (1992, August). A breeding trial using a transcervical technique for artificial insemination in sheep. In *Proc 12th Int Cong Anim Reprod* (Vol. 3, pp. 1531-1533).
- Buschbeck, C. (2020). About SRGenetics-WoolDrift Farm. Guelph System for Transcervical AI (GST-AI). The <https://wooldrift.com/srgenetics/about.htm>

343 Candappa, I. B., Bainbridge, H. C., Price, N. T., Hourigan, K. R., & Bartlewski, P. M.  
344 (2009). A preliminary study on the suitability of Cervidil® to induce cervical  
345 dilation for artificial insemination in ewes. *Research in veterinary*  
346 *science*, 87(2), 204-206.

347 Donovan, A., Hanrahan, J. P., Kummer, E., Duffy, P., & Boland, M. P. (2004). Fertility  
348 in the ewe following cervical insemination with fresh or frozen-thawed semen at  
349 a natural or synchronised oestrus. *Animal Reproduction Science*, 84(3-4), 359-  
350 368.

351 Donovan, A., Hanrahan, J. P., Lally, T., Boland, M., Byrne, G. P., Duffy, P., ... &  
352 O'Neill, D. J. (2001). *AI For Sheep Using Frozen-thawed Semen*. Teagasc.

353 Eppleston, J., Salamon, S., Moore, N. W., & Evans, G. (1994). The depth of cervical  
354 insemination and site of intrauterine insemination and their relationship to the  
355 fertility of frozen-thawed ram semen. *Animal Reproduction Science*, 36(3-4),  
356 211-225.

357 Evans, G., & Maxwell, W. C. (1987). *Salamons' artificial insemination of sheep and*  
358 *goats* (No. Ed. 2). Butterworths.

359 Faigl, V., Vass, N., Jávors, A., Kulcsár, M., Solti, L., Amiridis, G., & Cseh, S. (2012).  
360 Artificial insemination of small ruminants—A review. *Acta Veterinaria*  
361 *Hungarica*, 60(1), 115-129.

362 Falchi, L., Taema, M., La Clanche, S., & Scaramuzzi, R. J. (2012). The pattern of  
363 cervical penetration and the effect of topical treatment with prostaglandin and/or  
364 FSH and oxytocin on the depth of cervical penetration in the ewe during the  
365 peri-ovulatory period. *Theriogenology*, 78(2), 376-384.

366 Fernandez-Abella, D., Preve, M. O., & Villegas, N. (2003). Insemination time and  
367 dilution rate of cooled and chilled ram semen affects  
368 fertility. *Theriogenology*, 60(1), 21-26.

369 Fonseca, J. F., Souza, J. M. G., & Camargo, L. S. A. (2010). Produção de oócitos e  
370 embriões de pequenos ruminantes: passado, presente e futuro. *Acta Sci. Vet*, 38,  
371 337-369.

372 Fukui, Y., & Roberts, E. M. (1978). Further studies on non-surgical intrauterine  
373 technique for artificial insemination in the ewe. *Theriogenology*, 10(5), 381-393.

374 Gage, M. J. (1994). Associations between body size, mating pattern, testis size and  
375 sperm lengths across butterflies. *Proceedings of the Royal Society of London.*  
376 *Series B: Biological Sciences*, 258(1353), 247-254.

377 Gizaw, S., Tesfay, Y., Mekasha, Y., Mekuriaw, Z., Gugsu, T., Ebro, A., ... & Tegegne,  
378 A. (2016). Hormonal oestrus synchronization in four sheep breeds in Ethiopia:  
379 impacts on genetic improvement and flock productivity.

380 Gizaw, S., Tesfay, Y., Mekasha, Y., Mekuriaw, Z., Gugsu, T., Ebro, A., Gebremedhin,  
381 B., Hoekstra, D. and Tegegne, A. (2016). Hormonal oestrus synchronization in  
382 four sheep breeds in Ethiopia: impacts on genetic improvement and flock  
383 productivity. *LIVES Working Paper*.

384 Google Patents. 2012. Holding rack for artificial insemination of ewes-Google Patents.  
385 Retrieved from "<https://patents.google.com/patent/CN202497264U/en>">  
386 Abstract, p-1.

387 Halbert, G. W., Dobson, H. J. S. W., Walton, J. S., Sharpe, P., & Buckrell, B. C. (1990).  
388 Field evaluation of a technique for transcervical intrauterine insemination of  
389 ewes. *Theriogenology*, 33(6), 1231-1243.

390 Halbert, G. W., Dobson, H., Walton, J. S., & Buckrell, B. C. (1990C). A technique for  
391 transcervical intrauterine insemination of ewes. *Theriogenology*, 33(5), 993-  
392 1010.

393 Hawk, H. W. (1983). Sperm survival and transport in the female reproductive  
394 tract. *Journal of Dairy Science*, 66(12), 2645-2660.

395 Hernández Ballesteros, J. A., Navarrete Méndez, R., Benítez Meza, J. A., Moreno  
396 Flores, L. A., Gómez Gurrola, A., & Bernal Partida, M. A. (2015). Fertilidad  
397 con el uso de inseminación artificial en ovejas. *São Paulo: Editorial BM*.

398 Islam, R. (2011). Synchronization of estrus in cattle: A review. *Veterinary World*. 4 (3):  
399 136–141.

400 Jha, P. K., Alam, M. G. S., Mansur, A. A., Naher, N., Islam, T., Bhuiyan, M. U., &  
401 Bari, F. Y. (2019). Cryopreservation of Bangladeshi ram semen using different  
402 diluents and manual freezing techniques. *Cryobiology*, 89, 35-41.

403 Karagiannidis, A., Varsakeli, S., Karatzas, G., & Brozos, C. (2001). Effect of time of  
404 artificial insemination on fertility of progestagen and PMSG treated indigenous  
405 Greek ewes, during non-breeding season. *Small Ruminant Research*, 39(1), 67-  
406 71.

407 Khalifa, R. M., Sayre, B. L., & Lewis, G. S. (1992). Exogenous oxytocin dilates the  
408 cervix in ewes. *Journal of Animal Science*, 70(1), 38-42.

409 Kumar, D., & Naqvi, S. M. K. (2014). Effect of time and depth of insemination on  
410 fertility of Bharat Merino sheep inseminated trans-cervical with frozen-thawed  
411 semen. *Journal of Animal Science and Technology*, 56(1), 1-6.



- 412 Masoudi, R., Shahneh, A. Z., Towhidi, A., Kohram, H., Akbarisharif, A., Sharafi, M., ...  
 413 & Shahab-El-Deen, M. A. M. M. (2017). Cervical dilation and improvement of  
 414 reproductive performance in fat-tailed ewes via cervical dilator treatments. *Asian*  
 415 *Pacific Journal of Reproduction*, 6(2), 93.
- 416 Maxwell, W. M. C. (1980). Fertility of ram semen frozen in autumn and spring.  
 417 In *Australian Society for Reproductive Biology, Twelfth annual conference*,  
 418 *University of New England, Armidale, August 27-29, 1980. Program and*  
 419 *abstracts of papers..*
- 420 Maxwell, W. M. C., & Hewitt, L. J. (1986). A comparison of vaginal, cervical and  
 421 intrauterine insemination of sheep. *The Journal of Agricultural Science*, 106(1),  
 422 191-193.
- 423 Rekha, A., Zohara, B. F., Bari, F., & Alam, M. G. S. (2016). Comparison of commercial  
 424 Triladyl extender with a tris-fructose-egg-yolk extender on the quality of frozen  
 425 semen and pregnancy rate after transcervical AI in Bangladeshi indigenous  
 426 sheep (*Ovis aries*). *Small Ruminant Research*, 134, 39-43.
- 427 Salamon, S., & Maxwell, W. M. C. (1995). Frozen storage of ram semen I. Processing,  
 428 freezing, thawing and fertility after cervical insemination. *Animal Reproduction*  
 429 *Science*, 37(3-4), 185-249.
- 430 Sayre, B. L., & Lewis, G. S. (1996). Cervical dilation with exogenous oxytocin does not  
 431 affect sperm movement into the oviducts in ewes. *Theriogenology*, 45(8), 1523-  
 432 1533.
- 433 Schoenian, S. (2012). A beginner's guide to raising sheep. *University of Maryland*.  
 434 <http://www.sheep101.info/201/ewerepro.html>.
- 435 Smith, J. F., Parr, J., Beaumont, S., & Oliver, J. E. (1995). Comparison of cervical,  
 436 transcervical and laparoscopic insemination of ewes with chilled stored and  
 437 frozen ram semen. In *proceedings-New zealand society of animal*  
 438 *production* (Vol. 55, pp. 248-248). New zealand society of animal prod publ.
- 439 Stefanov, R., Krumova, E., Angelova, M., Dolashka, P., Voelter, W., & Zachariev, Z.  
 440 (2006). Artificial Insemination of Sheep and Cow with Semen Treated by  
 441 Cu/Zn-superoxide Dismutase from the Fungal Strain *Humicola lutea*  
 442 103. *World*, 1(1), 36-39.
- 443 Torrès, S., & Sevellec, C. (1987). Repeated superovulation and surgical recovery of  
 444 embryos in the ewe. *Reproduction Nutrition Développement*, 27(4), 859-863.

445 Tsuma, V. T., Khan, M. S., Okeyo Mwai, A., & Ibrahim, M. N. (2015). A training  
446 manual on artificial insemination in goats. ILRI Manual 19. Nairobi, Kenya:  
447 International Livestock Research Institute (ILRI).

448 Visser, D., & Salamon, S. (1974). Fertility following inseminations with frozen-thawed  
449 reconcentrated and unconcentrated ram semen. *Australian Journal of biological*  
450 *sciences*, 27(4), 423-426.

451 Wulster-Radcliffe, M. C., & Lewis, G. S. (2002). Development of a new transcervical  
452 artificial insemination method for sheep: effects of a new transcervical artificial  
453 insemination catheter and traversing the cervix on semen quality and  
454 fertility. *Theriogenology*, 58(7), 1361-1371.

455 Wulster-Radcliffe, M. C., Costine, B. A., & Lewis, G. S. (1999). Estradiol-17  $\beta$ -  
456 oxytocin-induced cervical dilation in sheep: application to transcervical embryo  
457 transfer. *Journal of animal science*, 77(10), 2587-2593.

458 Wulster-Radcliffe, M. C., Wang, S., & Lewis, G. S. (2004). Transcervical artificial  
459 insemination in sheep: effects of a new transcervical artificial insemination  
460 instrument and traversing the cervix on pregnancy and lambing  
461 rates. *Theriogenology*, 62(6), 990-1002.

462 Zohara, B. F., Md, M. F. I., Alamc, G. S., & Baric, F. Y. (2014). Comparison of estrus  
463 synchronization by PGF2 $\alpha$  and progestagen sponge with PMSG in indigenous  
464 ewes in Bangladesh. *GSTF Journal of Veterinary Science (JVet)*, 1(1).



**Figure 1. (A) Oestrus detection by aproned ram, (B) Traditional restrained TCAI, (C) Lab cradle restrained TCAI**

Table 1: Pregnancy rates in Bangladeshi indigenous sheep following different approaches of artificial insemination

Groups of animals	Approaches of TCAI	Pregnancy rate (%)	Chi-square test (P value)
Group 1 Restraining technique	Traditional	35 (7/20)	0.519
	Cradle	45 (9/20)	
Group 2 Time of TCAI	Single-time	35 (7/20)	0.204
	Double-times	55 (11/20)	
Group 3 Cervix ripening treatment	OT treatment	50 (5/10)	0.653
	OT + E2 treatment	55 (6/11)	

TCAI-trans-cervical artificial insemination, OT-Oxytocin, E2- Estradiol