

1 Original Article

2 **Analysis of the Ebola fractional-order model with Caputo-Fabrizio derivative**

3 Nita H Shah^{*}, Kapil Chaudhary

4 Department of Mathematics, Faculty of Science, Gujarat University,

5 Ahmedabad, 380009, India

6 ***Corresponding Author,**

7 **Email address: nitahshah@gmail.com, kapil.math.hons@gmail.com**

8
9 **Abstract**

10 This paper uses a fractional-order epidemic model to describe the transmission
11 dynamics of the Ebola virus. The proposed model uses the fractional-order derivative in
12 Caputo-Fabrizio's sense. It calculates the time-independent solutions of the proposed model,
13 and the next-generation matrix method is used to calculate the basic reproduction number. It
14 obtains the condition for the existence and uniqueness of the solutions of the model. Further,
15 the stability condition for generalized Ulam-Hyers-Rassias stability of the proposed model is
16 obtained. In numerical simulations, it shows how the proposed model's approximate solution
17 varies for integer and fractional orders. It also shows the behavior of the Ebola infections,
18 deceased and susceptible for various values of the contact rate. **To demonstrate efficiency**
19 **while using less time, CPU time is given in the tabular form.**

20
21 **Keywords:** Ebola, Caputo-Fabrizio derivative, Stability, Basic reproduction number,
22 Simulations

23

24 **2010 MSC:** 37Nxx, 37M05, 26A33

25 **1. Introduction**

26 The Filoviridae family member Ebola virus (EBOV) causes the inflammatory, severe,
27 potentially fatal disease known as (EVD) Ebola viral disease (used to be referred to as Ebola
28 hemorrhagic fever), which affects both humans and great apes. The first species of EBOV
29 was discovered near the Ebola River in the Democratic Republic of Congo in the Central
30 African continent in 1976 (Bisimwa, Biamba, Aborode, Cakwira, & Akilimali, 2022;
31 Feldmann, Sprecher, & Geisbert, 2020). With mortality rates reaching from 50% to 90% in
32 some instances, Death due to Ebola hemorrhagic fever can take place in as little as a few days
33 (Hammouch, Rasul, Ouakka, & Elazzouzi, 2022). The illness requires between two and
34 twenty-one days (typically, six to ten days) to incubate and eight hours to replicate (Feldmann
35 *et al.*, 2020). Humans can contract EBOV through close physical contact with infected bodily
36 fluids; blood, faeces, and vomit (World Health Organization [WHO], 2014). Over the past
37 three decades, EBOV has been responsible for a series of epidemics (Zhang & Jain, 2020).
38 The outbreak of 2013–16 was categorized by WHO as a Public Health Emergency of
39 International Concern, which highlighted the difficulties associated with treating Ebola virus
40 infections and raised concerns about society’s readiness to manage future epidemics on a
41 scientific, clinical, and sociological level.

42

43 **1.1.Literature Survey**

44 Many researchers have created epidemic models to better understand the EBOV virus
45 disease's mechanics. Some are integer order, and the rest are fractional-order models. **The**
46 **fractional-order model, in contrast to integer-order models, provides more freedom to fit the**

47 real data, which enhances the model's coherence with actual data and observations
48 (Khajehsaeid, 2018). In (Singh, 2020), the author used fractional-order GL-derivative to form
49 an iterative numerical scheme to find the numerical solutions of the epidemic model based
50 on EBOV and computed the CPU time usage. In (Srivastava, & Saad, 2020), the authors
51 looked at three potential kernel-based numerical solutions for the fractal-fractional Ebola
52 virus. In (Gao, Li, Li, & Zhou, 2021; Shaikh & Nisar, 2019), the authors used the fractional-
53 order CF-derivatives and fixed-point theorem in the proposed epidemic model to show the
54 existence and uniqueness of the governing system's solution. In (Shah, Patel, & Yeolekar,
55 2019), the authors proposed an integer-order model that discussed the vertical dynamics of
56 Ebola with media impacts. In (Liu, Fečkan, O'Regan, & Wang, 2019; Nwajeri, Oname, &
57 Onyenegecha, 2021), the authors derived the generalized UHR-stability results using
58 fractional-order CF- derivatives within their proposed models. Fractional calculus is
59 currently the focus of numerous studies of epidemic models. Multiple numerical and
60 analytical methods have been developed to solve fractional calculus problems. Some other
61 related models are discussed in (Hussain, Baleanu, & Adeel, 2020; Liu, Fečkan, & Wang,
62 2020; Solís-Pérez, Gómez-Aguilar, & Atangana, 2018). In (Singh, Srivastava, Hammouch,
63 & Nisar, 2021), the authors analyzed the stability conditions and the numerical results of the
64 proposed fractional-order model on COVID-19. In (Singh, Baleanu, Singh, & Dutta, 2021),
65 the authors looked at a non-integer order smoking model, utilized an iterative technique to
66 get numerical findings, and listed CPU time to illustrate the efficiency of solutions.

67

68 2. Preliminaries

69 **Definition 1.** *The fractional-order ϕ -derivative with $\phi \in (0,1]$ of function $f \in$*

70 $H^1[a, b]$ in Caputo's sense is defined as,

71
$${}^c D_t^\phi f(t) = \frac{1}{\Gamma(1-\phi)} \int_a^t f'(s)(t-s)^{-\phi} ds, \quad t > a \quad (1)$$

72 A new derivative is introduced using an exponential kernel to avoid the singularity at $t = s$
73 in **the** above expression (1).

74 **Definition 2.** (Losada & Nieto, 2015) The new fractional-order ϕ -derivative of a
75 function f in Caputo-Fabrizio's sense can be written as,

76
$${}^{CF} D_t^\phi f(t) = \frac{(2-\phi)M(\phi)}{2(1-\phi)} \int_a^t f'(s) \exp\left[-\frac{\phi(t-s)}{(1-\phi)}\right] ds, \quad t > a$$

77 where $M(\phi)$ is the normalizing constant function depending upon ϕ .

78 **Definition 3.** The Laplace transform of the fractional-order ϕ -derivative of a
79 function f in Caputo-Fabrizio's sense is defined as,

80
$$\mathcal{L}\{{}^{CF} D_t^\phi f(t), s\} = \frac{s\mathcal{L}\{f(t), s\} - f(0)}{(s + \phi(1-s))}, \quad s \geq 0.$$

81 **Definition 4.** The fractional integral of order ϕ of a function f in Caputo-Fabrizio's
82 sense is defined as,

83
$${}^{CF} J_t^\phi (f(t)) = \frac{2(1-\phi)}{(2-\phi)M(\phi)} f(t) + \frac{2\phi}{(2-\phi)M(\phi)} \int_a^t f(\tau) d\tau, \quad t > a.$$

84

85 3. Model formulation

86 The compartments of the model are defined as follows: Susceptible that are
87 uninfected (S), exposed to Ebola infection (E), infectious from the infection (I), Hospitalized
88 (H), Deceased or critically sick (D), and Recovered from infection (R). The total population
89 (N) is the sum of all the compartments. At any instance of time t ,

90
$$N(t) = S(t) + E(t) + I(t) + H(t) + D(t) + R(t)$$

91 The model has the following assumptions:

92 • Exposed individuals $E(t)$, infectious individuals ($I(t)$), Hospitalized individuals
 93 ($H(t)$), and Deceased or critically sick individuals ($D(t)$) are carriers of the EBOV at any
 94 instance of time t .

95 • Whenever any susceptible person (S) comes into touch with any deceased (D),
 96 hospitalized (H), exposed (E), or infectious (I) person, it acquires EBOV at the rate $\alpha_1(E +$
 97 $I + H + D)S/N$.

98 The meaning of the parameters used in the model is given in Table 1. The governing
 99 system of the fractional-order non-linear differential equations which describe the proposed
 100 epidemic model is as follows:

$$\begin{aligned}
 & {}^{CF}D_t^\phi E(t) = \frac{\alpha_1(E(t)+I(t)+D(t)+H(t))S(t)}{N(t)} - (\mu + \alpha_2)E(t) \\
 & {}^{CF}D_t^\phi I(t) = \alpha_2E(t) - (\alpha_3 + \alpha_4 + \mu)I(t) \\
 101 & {}^{CF}D_t^\phi H(t) = \alpha_3I(t) - (\alpha_5 + \alpha_6 + \mu)H(t) \\
 & {}^{CF}D_t^\phi D(t) = \alpha_4I(t) + \alpha_5H(t) - \mu D(t) \\
 & {}^{CF}D_t^\phi R(t) = \alpha_6H(t) - \mu R(t) \\
 & {}^{CF}D_t^\phi S(t) = B - \mu S(t) - \frac{\alpha_1(E(t)+I(t)+D(t)+H(t))S(t)}{N(t)}
 \end{aligned} \tag{2}$$

102 with non-negative initial condition, $(S(0), E(0), I(0), H(0), D(0), R(0)) \in \mathbb{R}_+^6$.

103 It can be rewritten in vector form as follows:

$$104 \quad {}^{CF}D_t^\phi \vec{\psi}(t) = \vec{\mathcal{K}}(t, E(t), I(t), H(t), D(t), R(t), S(t)) \tag{3}$$

105 for $\phi \in (0,1]$, $t \in J = [0, b]$ with following initial condition, $\tag{4}$

$$106 \quad \vec{\psi}(0) = \vec{\psi}_0 = (E(0), I(0), H(0), D(0), R(0), S(0))^T \tag{5}$$

107 where $\vec{\psi}(t) = (E(t), I(t), H(t), D(t), R(t), S(t))^T$

108 and $\vec{\mathcal{K}}(t) = (\mathcal{K}_1(t), \mathcal{K}_2(t), \mathcal{K}_3(t), \mathcal{K}_4(t), \mathcal{K}_5(t), \mathcal{K}_6(t))^T$.

109

110 4. Analysis of model

111 4.1. Equilibrium Points

112 In this section, the equilibrium points of system (2) are evaluated. The points are the
113 steady-state solution of the system (2). There are two equilibrium points of the proposed
114 system to be analyzed. The Ebola-free equilibrium point, E^0 is given by:

$$115 E^0 = \left\{ S = \frac{B}{\mu}, E = I = H = D = R = 0 \right\}$$

116 The endemic equilibrium point E^1 is given by:

$$117 E^1 = \left\{ S^* = \frac{A_1}{A_2}, E^* = \frac{A_3}{A_4}, I^* = \frac{A_5}{A_6}, H^* = \frac{A_7}{A_8}, D^* = \frac{A_9}{A_{10}}, R^* = \frac{A_{11}}{A_{12}} \right\}$$

118 where,

$$119 A_1 = N\mu(\mu + \alpha_5 + \alpha_6)(\mu + \alpha_3 + \alpha_4)(\mu + \alpha_2)$$

$$120 A_2 = \alpha_1(\mu^3 + (\alpha_2 + \alpha_3 + \alpha_4 + \alpha_5 + \alpha_6)\mu^2 + ((\alpha_3 + \alpha_4 + \alpha_5 + \alpha_6)\alpha_2 + (\alpha_5 + \alpha_6)(\alpha_3$$

121 $+ \alpha_4))\mu + ((\alpha_5 + \alpha_6)\alpha_4 + \alpha_3\alpha_5)\alpha_2)$

$$122 A_3 = -N\mu^5 - N(\alpha_2 + \alpha_3 + \alpha_4 + \alpha_5 + \alpha_6)\mu^4 + (-N(\alpha_3 + \alpha_4 + \alpha_5 + \alpha_6)\alpha_2 + B\alpha_1$$

123 $- N(\alpha_5 + \alpha_6)(\alpha_3 + \alpha_4))\mu^3 + (B\alpha_1 - N(\alpha_5 + \alpha_6)(\alpha_3 + \alpha_4))\alpha_2$
124 $+ B\alpha_1(\alpha_3 + \alpha_4 + \alpha_5 + \alpha_6))\mu^2 + B((\alpha_3 + \alpha_4 + \alpha_5 + \alpha_6)\alpha_2 + (\alpha_5$
125 $+ \alpha_6)(\alpha_3 + \alpha_4))\alpha_1\mu + ((\alpha_5 + \alpha_6)\alpha_4 + \alpha_3\alpha_5)B\alpha_2\alpha_1$

$$126 A_4 = (\mu + \alpha_2)(\mu^3 + (\alpha_2 + \alpha_3 + \alpha_4 + \alpha_5 + \alpha_6)\mu^2 + ((\alpha_3 + \alpha_4 + \alpha_5 + \alpha_6) + (\alpha_5$$

127 $+ \alpha_6)(\alpha_3 + \alpha_4))\mu + ((\alpha_5 + \alpha_6)\alpha_4 + \alpha_3\alpha_5)\alpha_2)\alpha_1$

$$\begin{aligned}
128 \quad A_5 &= (-N\mu^5 - N(\alpha_2 + \alpha_3 + \alpha_4 + \alpha_5 + \alpha_6)\mu^4 + (-N(\alpha_3 + \alpha_4 + \alpha_5 + \alpha_6)\alpha_2 - N(\alpha_5 \\
129 &\quad + \alpha_6)\alpha_4 - N(\alpha_5 + \alpha_6)\alpha_3 + B\alpha_1)\mu^3 + ((-N(\alpha_5 + \alpha_6)\alpha_4 - N(\alpha_5 + \alpha_6)\alpha_3 \\
130 &\quad + B\alpha_1)\alpha_2 + B\alpha_1(\alpha_3 + \alpha_4 + \alpha_5 + \alpha_6))\mu^2 + B((\alpha_3 + \alpha_4 + \alpha_5 + \alpha_6)\alpha_2 \\
131 &\quad + (\alpha_5 + \alpha_6)(\alpha_3 + \alpha_4))\alpha_1\mu + ((\alpha_5 + \alpha_6)\alpha_4 + \alpha_3\alpha_5)B\alpha_2\alpha_1)\alpha_2 \\
132 \quad A_6 &= (\mu + \alpha_2)(\mu + \alpha_3 + \alpha_4)(\mu^3 + (\alpha_2 + \alpha_3 + \alpha_4 + \alpha_5 + \alpha_6)\mu^2 + ((\alpha_3 + \alpha_4 + \alpha_5 \\
133 &\quad + \alpha_6)\alpha_2 + (\alpha_5 + \alpha_6)(\alpha_3 + \alpha_4))\mu + ((\alpha_5 + \alpha_6)\alpha_4 + \alpha_3\alpha_5)\alpha_2)\alpha_1 \\
134 \quad A_7 &= (-N\mu^5 - N(\alpha_2 + \alpha_3 + \alpha_4 + \alpha_5 + \alpha_6)\mu^4 + (-N(\alpha_3 + \alpha_4 + \alpha_5 + \alpha_6)\alpha_2 - N(\alpha_5 \\
135 &\quad + \alpha_6)\alpha_4 - N\alpha_3\alpha_5 - N\alpha_3\alpha_6 + B\alpha_1)\mu^3 + ((-N((\alpha_5 + \alpha_6)\alpha_4 + \alpha_3\alpha_5 \\
136 &\quad + \alpha_3\alpha_6) + B\alpha_1)\alpha_2 + B\alpha_1(\alpha_3 + \alpha_4 + \alpha_5 + \alpha_6))\mu^2 + B((\alpha_3 + \alpha_4 + \alpha_5 \\
137 &\quad + \alpha_6)\alpha_2 + (\alpha_5 + \alpha_6)(\alpha_3 + \alpha_4))\alpha_1\mu + ((\alpha_5 + \alpha_6)\alpha_4 + \alpha_5)B\alpha_2\alpha_1)\alpha_2\alpha_3 \\
138 \quad A_8 &= (\mu + \alpha_2)(\mu + \alpha_3 + \alpha_4)(\mu + \alpha_5 + \alpha_6)(\mu^3 + (\alpha_2 + \alpha_3 + \alpha_4 + \alpha_5 + \alpha_6)\mu^2 + ((\alpha_3 \\
139 &\quad + \alpha_4 + \alpha_5 + \alpha_6)\alpha_2 + (\alpha_5 + \alpha_6)(\alpha_3 + \alpha_4))\mu + ((\alpha_5 + \alpha_6)\alpha_4 \\
140 &\quad + \alpha_3\alpha_5)\alpha_2)\alpha_1 \\
141 \quad A_9 &= (-N\mu^5 - N(\alpha_2 + \alpha_3 + \alpha_4 + \alpha_5 + \alpha_6)\mu^4 + (-N(\alpha_3 + \alpha_4 + \alpha_5 + \alpha_6)\alpha_2 - N(\alpha_5 \\
142 &\quad + \alpha_6)\alpha_4 - N(\alpha_5 + \alpha_6)\alpha_3 + B\alpha_1)\mu^3 + ((-N(\alpha_5 + \alpha_6)\alpha_4 - N(\alpha_5 \\
143 &\quad + \alpha_6)\alpha_3 + B\alpha_1)\alpha_2 + B\alpha_1(\alpha_3 + \alpha_4 + \alpha_5 + \alpha_6))\mu^2 + B((\alpha_3 + \alpha_4 + \alpha_5 \\
144 &\quad + \alpha_6)\alpha_2 + (\alpha_5 + \alpha_6)(\alpha_3 + \alpha_4))\alpha_1\mu + ((\alpha_5 + \alpha_6)\alpha_4 + \alpha_5)B\alpha_2\alpha_1)\alpha_2(\alpha_4\mu \\
145 &\quad + (\alpha_5 + \alpha_6)\alpha_4 + \alpha_3\alpha_5 \\
146 \quad A_{10} &= \mu A_8
\end{aligned}$$

$$\begin{aligned}
147 \quad A_{11} &= (-N\mu^5 - N(\alpha_2 + \alpha_3 + \alpha_4 + \alpha_5 + \alpha_6)\mu^4 + (-N(\alpha_3 + \alpha_4 + \alpha_5 + \alpha_6)\alpha_2 - N(\alpha_5 \\
148 &\quad + \alpha_6)\alpha_4 - N\alpha_3\alpha_5 - N\alpha_3\alpha_6 + B\alpha_1)\mu^3 + ((-N(\alpha_5 + \alpha_6)\alpha_4 - N\alpha_3\alpha_5 \\
149 &\quad - N\alpha_3\alpha_6 + B\alpha_1)\alpha_2 + B\alpha_1(\alpha_3 + \alpha_4 + \alpha_5 + \alpha_6))\mu^2 + B((\alpha_3 + \alpha_4 + \alpha_5 \\
150 &\quad + \alpha_6)\alpha_2 + (\alpha_5 + \alpha_6)(\alpha_3 + \alpha_4))\alpha_1\mu + ((\alpha_5 + \alpha_6)\alpha_4 \\
151 &\quad + \alpha_3\alpha_5) \alpha_2\alpha_1 B)\alpha_2\alpha_6\alpha_3
\end{aligned}$$

$$\begin{aligned}
152 \quad A_{12} &= \mu(\mu + \alpha_2)(\mu + \alpha_3 + \alpha_4)(\mu + \alpha_5 + \alpha_6)(\mu^3 + (\alpha_2 + \alpha_3 + \alpha_4 + \alpha_5 + \alpha_6)\mu^2 + ((\alpha_3 \\
153 &\quad + \alpha_4 + \alpha_5 + \alpha_6)\alpha_2 + (\alpha_5 + \alpha_6)(\alpha_3 + \alpha_4))\mu + ((\alpha_5 + \alpha_6)\alpha_4 \\
154 &\quad + \alpha_3\alpha_5)\alpha_2)\alpha_1
\end{aligned}$$

155

156 4.2 Basic Reproduction Number

157 The next-generation matrix method (Diekmann, Heesterbeek, & Metz, 1990;
158 Otunuga, 2021) is used to calculate the basic reproduction number of the proposed model.
159 The fractional-order system (2) can be rewritten as:

$$160 \quad {}^C D_t^\phi \vec{\psi}(t) = \vec{f}(t) - \vec{v}(t)$$

161 where,

$$162 \quad \vec{\psi}(t) = \begin{bmatrix} E(t) \\ I(t) \\ H(t) \\ D(t) \\ R(t) \\ S(t) \end{bmatrix}, \vec{f}(t) = \begin{bmatrix} \frac{\alpha_1(E + I + D + H)S}{N} \\ 0 \\ 0 \\ 0 \\ 0 \\ -\frac{\alpha_1(E + I + D + H)S}{N} \end{bmatrix} \text{ and } \vec{v}(t) = \begin{bmatrix} (\mu + \alpha_2)E \\ (\mu + \alpha_3 + \alpha_4)I - \alpha_2E \\ (\mu + \alpha_5 + \alpha_6)H - \alpha_3I \\ \mu D - \alpha_4I - \alpha_5H \\ \mu R - \alpha_6H \\ \mu S - B \end{bmatrix}$$

163 are the column vectors with the initial condition, $\vec{\psi}(0) = \vec{\psi}_0$. Assume that F and V be the
164 Jacobian matrices of the column vectors \vec{f} and \vec{v} , respectively.

165 At the Ebola-free equilibrium point E^0 ,

$$166 \quad F_{E^0} = F(E^0) = \begin{bmatrix} v & v & v & v & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 \\ -v & -v & -v & -v & 0 & 0 \end{bmatrix} \quad (\text{where } v = B\alpha_1 N\mu)$$

$$167 \quad V_{E^0} = V(E^0) = \begin{bmatrix} \mu + \alpha_2 & 0 & 0 & 0 & 0 & 0 \\ -\alpha_2 & \mu + \alpha_3 + \alpha_4 & 0 & 0 & 0 & 0 \\ 0 & -\alpha_3 & \mu + \alpha_5 + \alpha_6 & 0 & 0 & 0 \\ 0 & -\alpha_4 & -\alpha_5 & \mu & 0 & 0 \\ 0 & 0 & -\alpha_6 & 0 & \mu & 0 \\ 0 & 0 & 0 & 0 & 0 & \mu \end{bmatrix}$$

168 are the singular and non-singular 6×6 matrices. The basic reproduction number is the
169 spectral radius of the matrix FV^{-1} at Ebola-free point E^0 .

$$170 \quad \mathcal{R}_0 = \rho(F_{E^0}V_{E^0}^{-1}) = \left[\frac{B(\mu^3 + (\alpha_2 + \alpha_3 + \alpha_4 + \alpha_5 + \alpha_6)\mu^2 + ((\alpha_3 + \alpha_4 + \alpha_5 + \alpha_6)\alpha_2 + (\alpha_5 + \alpha_6)(\alpha_3 + \alpha_4))\mu + ((\alpha_5 + \alpha_6)\alpha_4 + \alpha_3\alpha_5)\alpha_2)\alpha_1}{N(\mu + \alpha_5 + \alpha_6)(\mu + \alpha_3 + \alpha_4)(\mu + \alpha_2)\mu^2} \right]$$

171

172 4.3 Invariant positive region

173 **Lemma 1.** *The proposed fractional-order model for Ebola infection (2)'s the feasible*
174 *domain of solution is*

$$175 \quad \Omega = \left\{ (S, E, I, H, D, R) \in \mathbb{R}_+^6 \mid 0 \leq N \leq \frac{B}{\mu} \right\}$$

176 *positively invariant.*

177 *Proof.* Adding all the equations in the fractional-order system (2),

$$178 \quad {}^{CF}D_t^\phi N(t) = B - \mu N(t)$$

179 Using the Laplace transform for both sides of the previous equation,

$$180 \quad \mathcal{L}\{ {}^{CF}D_t^\phi N(t), s \} = \mathcal{L}\{ B - \mu N(t), s \}$$

181 This gives,

$$182 \quad \frac{s\mathcal{L}\{N(t), s\} - N(0)}{(s + \phi(1 - s))} = \frac{B}{s} - \mu\mathcal{L}\{N(t), s\}$$

183 provided $s > 0$.

184 Solving for $\mathcal{L}\{N(t), s\}$ and taking inverse Laplace transform,

$$185 \quad N(t) = N(0)\mathcal{L}^{-1}\left\{\frac{1}{s + \mu s + \mu\phi(1 - s)}\right\} + B\mathcal{L}^{-1}\left\{\frac{s + \phi(1 - s)}{(1 + \mu - \phi\mu)s^2 + \phi\mu s}\right\}$$

$$186 \quad = \frac{N(0)}{1 + \mu - \mu\phi} \mathcal{L}^{-1}\left\{\frac{1}{s + \left(\frac{\mu\phi}{1 + \mu(1 - \phi)}\right)}\right\} + \frac{B}{1 + \mu - \mu\phi} \left[(1 - \phi)\mathcal{L}^{-1}\left\{\frac{1}{s + \frac{\mu\phi}{1 + \mu(1 - \phi)}}\right\} \right. \\ \left. + \phi\mathcal{L}^{-1}\left\{\frac{1}{s\left(s + \frac{\phi\mu}{(1 + \mu - \mu\phi)}\right)}\right\} \right]$$

$$187 \quad = \frac{N(0)}{1 + \mu - \mu\phi} \left[\exp\left[-\left(\frac{\mu\phi t}{1 + \mu(1 - \phi)}\right)\right] \right]$$

$$188 \quad + \frac{B}{1 + \mu - \mu\phi} \left[(1 - \phi)\exp\left[-\left(\frac{\mu\phi t}{1 + \mu(1 - \phi)}\right)\right] \right. \\ \left. + \frac{1 + \mu - \mu\phi}{\mu} \left[1 - \exp\left[-\left(\frac{\mu\phi t}{1 + \mu(1 - \phi)}\right)\right] \right] \right]$$

$$189 \quad = \frac{B}{\mu} + \left[\frac{N(0) + B(1 - \phi)}{1 + \mu - \mu\phi} - \frac{B}{\mu} \right] \exp\left[-\left(\frac{\mu\phi t}{1 + \mu(1 - \phi)}\right)\right]$$

190 and because of the asymptotic decay characteristic of **the** inverse exponential function as time

191 grows, $\lim_{t \rightarrow \infty} N(t) \leq B/\mu$. Therefore, the fractional-order system has a positively oriented

192 bounded region.

193

194 **5. Existence and Uniqueness of the solution**

195 In this section, the existence and uniqueness of the solution of the proposed fractional-

196 order model are shown. Applying the fractional-integral operator (${}^{CF}J_t^\phi$) on both sides into

197 the system of equations (2),

$$\begin{aligned}
E(t) - E(0) &= {}^{CF} J_t^\phi \left(\frac{\alpha_1(E(t)+I(t)+D(t)+H(t))S(t)}{N(t)} - (\mu + \alpha_2)E(t) \right) \\
I(t) - I(0) &= {}^{CF} J_t^\phi (\alpha_2 E(t) - (\alpha_3 + \alpha_4 + \mu)I(t)) \\
H(t) - H(0) &= {}^{CF} J_t^\phi (\alpha_3 I(t) - (\alpha_5 + \alpha_6 + \mu)H(t)) \\
D(t) - D(0) &= {}^{CF} J_t^\phi (\alpha_4 I(t) + \alpha_5 H(t) - \mu D(t)) \\
R(t) - R(0) &= {}^{CF} J_t^\phi (\alpha_6 H(t) - \mu R(t)) \\
S(t) - S(0) &= {}^{CF} J_t^\phi \left(B - \mu S(t) - \frac{\alpha_1(E(t)+I(t)+D(t)+H(t))S(t)}{N(t)} \right)
\end{aligned} \tag{6}$$

199 Solving the right-hand side of the system (6),

$$\begin{aligned}
E(t) - E(0) &= \frac{2(1-\phi)}{(2-\phi)M(\phi)} \mathcal{K}_1(t, \psi(t)) + \frac{2\phi}{(2-\phi)M(\phi)} \int_0^t \mathcal{K}_1(\tau, \psi(\tau)) d\tau \\
I(t) - I(0) &= \frac{2(1-\phi)}{(2-\phi)M(\phi)} \mathcal{K}_2(t, \psi(t)) + \frac{2\phi}{(2-\phi)M(\phi)} \int_0^t \mathcal{K}_2(\tau, \psi(\tau)) d\tau \\
H(t) - H(0) &= \frac{2(1-\phi)}{(2-\phi)M(\phi)} \mathcal{K}_3(t, \psi(t)) + \frac{2\phi}{(2-\phi)M(\phi)} \int_0^t \mathcal{K}_3(\tau, \psi(\tau)) d\tau \\
D(t) - D(0) &= \frac{2(1-\phi)}{(2-\phi)M(\phi)} \mathcal{K}_4(t, \psi(t)) + \frac{2\phi}{(2-\phi)M(\phi)} \int_0^t \mathcal{K}_4(\tau, \psi(\tau)) d\tau \\
R(t) - R(0) &= \frac{2(1-\phi)}{(2-\phi)M(\phi)} \mathcal{K}_5(t, \psi(t)) + \frac{2\phi}{(2-\phi)M(\phi)} \int_0^t \mathcal{K}_5(\tau, \psi(\tau)) d\tau \\
S(t) - S(0) &= \frac{2(1-\phi)}{(2-\phi)M(\phi)} \mathcal{K}_6(t, \psi(t)) + \frac{2\phi}{(2-\phi)M(\phi)} \int_0^t \mathcal{K}_6(\tau, \psi(\tau)) d\tau
\end{aligned} \tag{7}$$

201 where kernels $\mathcal{K}_1, \mathcal{K}_2, \mathcal{K}_3, \mathcal{K}_4, \mathcal{K}_5,$ and \mathcal{K}_6 are defined as:

$$\begin{aligned}
\mathcal{K}_1(t, \psi(t)) &= \frac{\alpha_1(E(t)+I(t)+D(t)+H(t))S(t)}{N(t)} - (\mu + \alpha_2)E(t) \\
\mathcal{K}_2(t, \psi(t)) &= \alpha_2 E(t) - (\alpha_3 + \alpha_4 + \mu)I(t) \\
\mathcal{K}_3(t, \psi(t)) &= \alpha_3 I(t) - (\alpha_5 + \alpha_6 + \mu)H(t) \\
\mathcal{K}_4(t, \psi(t)) &= \alpha_4 I(t) + \alpha_5 H(t) - \mu D(t) \\
\mathcal{K}_5(t, \psi(t)) &= \alpha_6 H(t) - \mu R(t) \\
\mathcal{K}_6(t, \psi(t)) &= B - \mu S(t) - \frac{\alpha_1(E(t)+I(t)+D(t)+H(t))S(t)}{N(t)}
\end{aligned} \tag{8}$$

203 **Lemma 2.** The kernels $\mathcal{K}_1, \mathcal{K}_2, \mathcal{K}_3, \mathcal{K}_4, \mathcal{K}_5,$ and \mathcal{K}_6 satisfy Lipschitz's condition if

$$204 \quad 0 \leq L = \sup\{L_1, L_2, L_3, L_4, L_5, L_6\} < 1$$

205 where $L_1 = |\alpha_1 s_1 - (\mu + \alpha_2)|, L_2 = |\mu + \alpha_3 + \alpha_4|, L_3 = |\mu + \alpha_5 + \alpha_6|, L_4 = L_5 = \mu, L_6 =$

206 $|\alpha_1(1 - s_0 - r_0) - \mu|, s_0 = \inf_t S(t)/N(t) \leq \sup_t S(t)/N(t) = s_1$ and $r_0 = \inf_t R(t)/$

207 $N(t).$

208 *Proof.* Let E_1, E_2 be corresponding functions for the kernel \mathcal{K}_1 . Let I_1, I_2 be corresponding
209 functions for the kernel \mathcal{K}_2 . Let H_1, H_2 be corresponding functions for the kernel \mathcal{K}_3 . Let
210 D_1, D_2 be corresponding functions for the kernel \mathcal{K}_4 , R_1, R_2 be corresponding functions for
211 the kernel \mathcal{K}_5 and S_1, S_2 be corresponding functions for the kernel \mathcal{K}_6 . Then,

$$212 \quad \|\mathcal{K}_1(t, E_1(t)) - \mathcal{K}_1(t, E_2(t))\| = \left\| \left(\frac{\alpha_1 S(t)}{N(t)} - (\mu + \alpha_2) \right) (E_1(t) - E_2(t)) \right\| \quad (9)$$

$$213 \quad \leq \underbrace{\sup_t \left| \frac{\alpha_1 S(t)}{N(t)} - (\mu + \alpha_2) \right|}_{L_1} \|E_1(t) - E_2(t)\| \quad (10)$$

214 Similarly,

$$215 \quad \|\mathcal{K}_2(t, I_1(t)) - \mathcal{K}_2(t, I_2(t))\| = \|(-(\mu + \alpha_3 + \alpha_4))(I_1(t) - I_2(t))\| \quad (11)$$

$$216 \quad \leq \underbrace{|\mu + \alpha_3 + \alpha_4|}_{L_2} \|I_1(t) - I_2(t)\| \quad (12)$$

$$217 \quad \|\mathcal{K}_3(t, H_1(t)) - \mathcal{K}_3(t, H_2(t))\| = \|(-(\mu + \alpha_5 + \alpha_6))(H_1(t) - H_2(t))\| \quad (13)$$

$$218 \quad \leq \underbrace{|\mu + \alpha_5 + \alpha_6|}_{L_3} \|H_1(t) - H_2(t)\| \quad (14)$$

$$219 \quad \|\mathcal{K}_4(t, D_1(t)) - \mathcal{K}_3(t, D_2(t))\| = \|(-\mu)(D_1(t) - D_2(t))\| \quad (15)$$

$$220 \quad \leq \underbrace{|\mu|}_{L_4} \|D_1(t) - D_2(t)\| \quad (16)$$

$$221 \quad \|\mathcal{K}_5(t, R_1(t)) - \mathcal{K}_5(t, R_2(t))\| = \|(-\mu)(R_1(t) - R_2(t))\| \quad (17)$$

$$222 \quad \leq \underbrace{|\mu|}_{L_5} \|R_1(t) - R_2(t)\| \quad (18)$$

$$223 \quad \|\mathcal{K}_6(t, S_1(t)) - \mathcal{K}_6(t, S_2(t))\| = \left\| \frac{\alpha_1(E(t)+I(t)+D(t)+H(t))(S_1(t)-S_2(t))}{N(t)} \right\| \quad (19)$$

$$224 \quad \leq \underbrace{\sup_t \left| \alpha_1 \left(1 - \frac{S(t)}{N(t)} - \frac{R(t)}{N(t)} \right) \right|}_{L_6} \|S_1(t) - S_2(t)\| \quad (20)$$

225 For each $n \in \mathbb{N}$, we can get the following system of recursive relations using Picard's
226 iteration,

$$\begin{aligned}
227 \quad E_n(t) &= \frac{2(1-\phi)}{(2-\phi)M(\phi)} \mathcal{K}_1(t, E_{n-1}(t)) + \frac{2\phi}{(2-\phi)M(\phi)} \int_0^t \mathcal{K}_1(\tau, E_{n-1}(\tau)) d\tau \\
228 \quad I_n(t) &= \frac{2(1-\phi)}{(2-\phi)M(\phi)} \mathcal{K}_2(t, I_{n-1}(t)) + \frac{2\phi}{(2-\phi)M(\phi)} \int_0^t \mathcal{K}_2(\tau, I_{n-1}(\tau)) d\tau \\
229 \quad H_n(t) &= \frac{2(1-\phi)}{(2-\phi)M(\phi)} \mathcal{K}_3(t, H_{n-1}(t)) + \frac{2\phi}{(2-\phi)M(\phi)} \int_0^t \mathcal{K}_3(\tau, H_{n-1}(\tau)) d\tau \\
230 \quad D_n(t) &= \frac{2(1-\phi)}{(2-\phi)M(\phi)} \mathcal{K}_4(t, D_{n-1}(t)) + \frac{2\phi}{(2-\phi)M(\phi)} \int_0^t \mathcal{K}_4(\tau, D_{n-1}(\tau)) d\tau \\
231 \quad R_n(t) &= \frac{2(1-\phi)}{(2-\phi)M(\phi)} \mathcal{K}_5(t, R_{n-1}(t)) + \frac{2\phi}{(2-\phi)M(\phi)} \int_0^t \mathcal{K}_5(\tau, R_{n-1}(\tau)) d\tau \\
232 \quad S_n(t) &= \frac{2(1-\phi)}{(2-\phi)M(\phi)} \mathcal{K}_6(t, S_{n-1}(t)) + \frac{2\phi}{(2-\phi)M(\phi)} \int_0^t \mathcal{K}_6(\tau, S_{n-1}(\tau)) d\tau \quad (21)
\end{aligned}$$

233 Now, using Lipschitz's inequalities and above recursive equations (21),

$$234 \quad \|\Delta E_n(t)\| = \|E_n(t) - E_{n-1}(t)\| \quad (22)$$

$$235 \quad \leq \frac{2(1-\phi)L_1}{(2-\phi)M(\phi)} \|\Delta E_{n-1}(t)\| + \frac{2\phi L_1}{(2-\phi)M(\phi)} \int_0^t \|\Delta E_{n-1}(\tau)\| d\tau \quad (23)$$

$$236 \quad \leq \left(\frac{2L_1(1-\phi+\phi T_{\text{sup}})}{M(\phi)(2-\phi)} \right) \|\Delta E_{n-1}(t)\| \quad (24)$$

237 Similarly,

$$238 \quad \|\Delta I_n(t)\| = \|I_n(t) - I_{n-1}(t)\| \leq \left(\frac{2L_2(1-\phi+\phi T_{\text{sup}})}{M(\phi)(2-\phi)} \right) \|\Delta I_{n-1}(t)\| \quad (25)$$

$$239 \quad \|\Delta H_n(t)\| = \|H_n(t) - H_{n-1}(t)\| \leq \left(\frac{2L_3(1-\phi+\phi T_{\text{sup}})}{M(\phi)(2-\phi)} \right) \|\Delta H_{n-1}(t)\| \quad (26)$$

$$240 \quad \|\Delta D_n(t)\| = \|D_n(t) - D_{n-1}(t)\| \leq \left(\frac{2L_4(1-\phi+\phi T_{\text{sup}})}{M(\phi)(2-\phi)} \right) \|\Delta D_{n-1}(t)\| \quad (27)$$

$$241 \quad \|\Delta R_n(t)\| = \|R_n(t) - R_{n-1}(t)\| \leq \left(\frac{2L_5(1-\phi+\phi T_{\text{sup}})}{M(\phi)(2-\phi)} \right) \|\Delta R_{n-1}(t)\| \quad (28)$$

$$242 \quad \|\Delta S_n(t)\| = \|S_n(t) - S_{n-1}(t)\| \leq \left(\frac{2L_6(1-\phi+\phi T_{\text{sup}})}{M(\phi)(2-\phi)} \right) \|\Delta S_{n-1}(t)\| \quad (29)$$

243 Further, it can be observed using telescopic sum that,

$$244 \quad E_m(t) = E_0 + \sum_{n=1}^m \Delta E_n(t), \quad I_m(t) = I_0 + \sum_{n=1}^m \Delta I_n(t)$$

$$245 \quad H_m(t) = H_0 + \sum_{n=1}^m \Delta H_n(t) \quad D_m(t) = D_0 + \sum_{n=1}^m \Delta D_n(t)$$

246
$$R_m(t) = R_0 + \sum_{n=1}^m \Delta R_n(t), S_m(t) = S_0 + \sum_{n=1}^m \Delta S_n(t)$$

247 This proves the result.

248 **Theorem 1. (Existence of solution)** *There exists a solution of the fractional system*

249 *(2) provided if $0 \leq \theta < 1$. where $\theta = \left(\frac{2L(1-\phi+\phi T_{sup})}{M(\phi)(2-\phi)}\right)$ and $L = \sup\{L_1, L_2, L_3, L_4, L_5, L_6\}$.*

250 *Proof.* The functions $E(t), I(t), H(t), D(t), R(t), S(t)$ are bounded and respective kernels

251 satisfy the Lipschitz's conditions. Using the recursive formula for the inequalities (22) - (29),

252
$$\| \Delta E_n(t) \| \leq \left(\frac{2L_1(1-\phi+\phi T_{sup})}{M(\phi)(2-\phi)}\right)^{n-1} \| \Delta E_1(t) \|$$

253
$$\leq \theta^{n-1} \| \Delta E_1(t) \|$$

254
$$\xrightarrow{n \rightarrow \infty} 0, \text{ as } 0 \leq \theta < 1.$$

255 Similarly, it can be observed that each of the following sequences of,

256
$$\| \Delta I_n(t) \|, \| \Delta H_n(t) \|, \| \Delta D_n(t) \|, \| \Delta R_n(t) \|, \| \Delta S_n(t) \| \xrightarrow{n \rightarrow \infty} 0, \text{ as } 0 \leq \theta < 1.$$

257 This proves the solutions of the fractional system exist and are of the form mentioned in (7).

258 **Lemma 3.** (Nwajeri, Panle, Omame, Obi, & Onyenegecha, 2022) *Consider the initial*

259 *value problem ${}^{CF}\psi_t^\phi(t) = \mathcal{K}(t, \psi(t)), \psi(0) = \psi_0$ and suppose that there exists a*

260 *Lipschitz's constant $L \geq 0$ such that*

261
$$|\mathcal{K}(t, \psi_1(t)) - \mathcal{K}(t, \psi_2(t))| \leq L|\psi_1(t) - \psi_2(t)|, \quad (30)$$

262 *for all $t \in J = [0, b]$ and $\psi_1, \psi_2 \in C(J, \mathbb{R})$. If $L \left(\frac{2(1-\phi)+2\phi T_{sup}}{(2-\phi)M(\phi)}\right) < 1$, Then, there exists a*

263 *unique solution of the initial value problem on $J = [0, b]$.*

264 *Proof.* The uniqueness of the solution to this initial value problem is the consequence of the

265 Banach-Fixed point theorem. Let $C(J, \mathbb{R})$ denotes the Banach space of all the continuous

266 functions from J to \mathbb{R} with infinity-norm.

267 $\|f\|_\infty = \sup_t \{|f(t)| | t \in J = [0, b]\}, \quad \forall f \in C(J, \mathbb{R})$

268 Consider the mapping $\omega: C^\theta(J, \mathbb{R}) \rightarrow C^\theta(J, \mathbb{R})$ defined by,

269
$$\omega\psi(t) = \psi_0 + \frac{2(1-\phi)}{(2-\phi)M(\phi)} (\mathcal{K}(t) - \mathcal{K}(0)) + \frac{2\phi}{(2-\phi)M(\phi)} \int_0^t \mathcal{K}(\tau) d\tau$$

270 Assume $\psi_1, \psi_2 \in C^\theta(J, \mathbb{R})$ and for each $t \in J$,

271
$$|\omega\psi_1(t) - \omega\psi_2(t)| \leq \frac{2(1-\phi)}{(2-\phi)M(\phi)} |\mathcal{K}(t, \psi_1(t)) - \mathcal{K}(t, \psi_2(t))|$$

272
$$+ \frac{2\phi}{(2-\phi)M(\phi)} \int_0^t |\mathcal{K}(\tau, \psi_1(\tau)) - \mathcal{K}(\tau, \psi_2(\tau))| d\tau$$

273 Using inequality (30) and for each $t \in J$,

274
$$|\omega\psi_1(t) - \omega\psi_2(t)| \leq \frac{2L(1-\phi)}{(2-\phi)M(\phi)} |\psi_1(t) - \psi_2(t)|$$

275
$$+ \frac{2L\phi}{(2-\phi)M(\phi)} \int_0^t |\psi_1(\tau) - \psi_2(\tau)| d\tau$$

276
$$\leq L \left(\frac{2(1-\phi)}{(2-\phi)M(\phi)} + \frac{2\phi T_{\text{sup}}}{(2-\phi)M(\phi)} \right) |\psi_1(t) - \psi_2(t)|$$

277 Taking supremum over $t \in J$,

278
$$\|\omega\psi_1 - \omega\psi_2\|_\infty \leq L \left(\frac{2(1-\phi) + 2\phi T_{\text{sup}}}{(2-\phi)M(\phi)} \right) \|\psi_1 - \psi_2\|_\infty$$

279 Thus, ω is a contraction mapping if $0 \leq L \left(\frac{2(1-\phi) + 2\phi T_{\text{sup}}}{(2-\phi)M(\phi)} \right) < 1$. Consequently, by Banach-

280 fixed point theorem, the operator ω has a fixed point say (ψ) i.e. $(\omega\psi = \psi)$ which is the

281 required unique solution of the initial value problem on $C(J, \mathbb{R})$.

282 **Theorem 2. (Uniqueness of solution)** *The solution (as mentioned in system 7) of the*

283 *fractional system (2) is unique.*

284 *Proof.* We use Lemma 3 to show the uniqueness of the solution (as mentioned in system 7)

285 for the fractional system (2). By Lemma 2, kernels $\mathcal{K}_1, \mathcal{K}_2, \mathcal{K}_3, \mathcal{K}_4, \mathcal{K}_5$, and \mathcal{K}_6 satisfies the

286 Lipschitz's conditions with constants L_1, L_2, L_3, L_4, L_5 , and L_6 , respectively. Letting $L =$

287 $\sup\{L_1, L_2, L_3, L_4, L_5, L_6\}$ and $\theta = \left(\frac{2(1-\phi)+2\phi T_{\text{sup}}}{(2-\phi)M(\phi)}\right)$. Since solution exists if $0 \leq \theta < 1$
 288 (hypothesis of existence Theorem 1). Thus, the hypothesis of the Lemma 3 is satisfied for
 289 each of the equations (with initial values) of the fractional-order system (2). Hence, using
 290 Lemma 3, we conclude that the fractional-order system (2) has a unique solution if $0 \leq \theta <$
 291 1.

292

293 6. Stability

294 This section obtains the appropriate stability condition for the generalized Ulam-
 295 Hyers-Rassias stability of the proposed fractional-order model.

296

297 6.1. Generalized Ulam-Hyers-Rassias stability

298 **Definition 5.** (Nwajeri *et al.*, 2021, 2022; Liu, Fečkan, O'Regan, & Wang, 2019) *The*
 299 *fractional-order model* ${}^{CF}D_t^\phi \psi(t) = \mathcal{K}(t, \psi(t))$ *is generalized Ulam-Hyers-Rassias*
 300 *(UHR) stable in accordance with* $\mathcal{Y}(t) \in H^1[J, \mathbb{R}^+]$ *if there exists a positive real value* ε_ϕ
 301 *(depending upon ϕ) such that for every solution ψ of the following inequality,*

$$302 \quad |{}^{CF}D_t^\phi \psi(t) - \mathcal{K}(t, \psi(t))| \leq \mathcal{Y}(t),$$

303 *There exists a solution* $\tilde{\psi} \in H^1(J, \mathbb{R}^+)$ *of the model with the following,*

$$304 \quad |\psi(t) - \tilde{\psi}(t)| \leq \varepsilon_\phi \mathcal{Y}(t) \quad \text{for each } t \in J.$$

305 **Lemma 4.** *The fractional-order model* ${}^{CF}D_t^\phi \psi(t) = \mathcal{K}(t, \psi(t))$ *(satisfying*
 306 *Lipschitz's condition with Lipschitz constant* L *depending upon kernel* \mathcal{K}) *is generalized*
 307 *UHR-stable in accordance with non-decreasing positive function* \mathcal{Y} *if,*

$$308 \quad 0 \leq \theta = \frac{2L((1-\phi)+\phi T_{\text{sup}})}{(2-\phi)M(\phi)} < 1 \quad (31)$$

309 *Proof.* We let $\mathcal{Y}(t)$ represent any arbitrary positive function, then there exists a positive real
 310 number η such that,

$$311 \quad \left(2(1 - \phi)\mathcal{Y}(t) + 2\phi \int_0^t \mathcal{Y}(\tau)d\tau\right) \leq \eta\mathcal{Y}(t). \quad (32)$$

312 Since the kernel of the fractional-order model satisfies Lipschitz's condition with Lipschitz's
 313 constant L (depending upon kernel \mathcal{K}) i.e.

$$314 \quad |\mathcal{K}(t, \psi(t)) - \mathcal{K}(t, \tilde{\psi}(t))| \leq L|\psi(t) - \tilde{\psi}(t)| \quad (33)$$

315 So, using the existence and uniqueness theorem of the model, there exists a unique solution
 316 say ($\tilde{\psi}$) of the fractional-order model of the following form,

$$317 \quad \tilde{\psi}(t) = \tilde{\psi}_0 + \frac{2(1-\phi)}{(2-\phi)M(\phi)}\mathcal{K}(t, \tilde{\psi}(t)) + \frac{2\phi}{(2-\phi)M(\phi)}\int_0^t \mathcal{K}(\tau, \tilde{\psi}(\tau))d\tau \quad (34)$$

318 Assume that ψ is the solution of the following inequality,

$$319 \quad |{}^{CF}D_t^\phi \psi(t) - \mathcal{K}(t, \psi(t))| \leq \mathcal{Y}(t), \quad (35)$$

320 Applying fractional-order integral operator,

$$321 \quad |\psi(t) - {}^{CF}J_t^\phi \mathcal{K}(t, \psi(t))| \leq \frac{\eta\mathcal{Y}(t)}{(2-\phi)M(\phi)} \quad (36)$$

$$322 \quad \left| \psi(t) - \tilde{\psi}_0 - \frac{2(1-\phi)}{(2-\phi)M(\phi)}\mathcal{K}(t, \psi(t)) - \frac{2\phi}{(2-\phi)M(\phi)}\int_0^t \mathcal{K}(\tau, \psi(\tau))d\tau \right| \leq \frac{\eta\mathcal{Y}(t)}{(2-\phi)M(\phi)} \quad (37)$$

323 Now, Consider the following,

$$324 \quad |\psi(t) - \tilde{\psi}(t)|$$

$$325 \quad = \left| \psi(t) - \tilde{\psi}_0 - \frac{2(1-\phi)}{(2-\phi)M(\phi)}\mathcal{K}(t, \tilde{\psi}(t)) - \frac{2\phi}{(2-\phi)M(\phi)}\int_0^t \mathcal{K}(\tau, \tilde{\psi}(\tau))d\tau \right|$$

$$326 \quad = \left| \begin{aligned} &\psi(t) - \tilde{\psi}_0 - \frac{2(1-\phi)}{(2-\phi)M(\phi)}\mathcal{K}(t, \tilde{\psi}(t)) - \frac{2\phi}{(2-\phi)M(\phi)}\int_0^t \mathcal{K}(\tau, \tilde{\psi}(\tau))d\tau \\ &+ \frac{2(1-\phi)}{(2-\phi)M(\phi)}\mathcal{K}(t, \psi(t)) + \frac{2\phi}{(2-\phi)M(\phi)}\int_0^t \mathcal{K}(\tau, \psi(\tau))d\tau \\ &- \frac{2(1-\phi)}{(2-\phi)M(\phi)}\mathcal{K}(t, \psi(t)) - \frac{2\phi}{(2-\phi)M(\phi)}\int_0^t \mathcal{K}(\tau, \psi(\tau))d\tau \end{aligned} \right|$$

$$\begin{aligned}
327 \quad & \leq \left| \psi(t) - \tilde{\psi}_0 - \frac{2(1-\phi)}{(2-\phi)M(\phi)} \mathcal{K}(t, \psi(t)) - \frac{2\phi}{(2-\phi)M(\phi)} \int_0^t \mathcal{K}(\tau, \psi(\tau)) d\tau \right| \\
328 \quad & \quad + \frac{2(1-\phi)}{(2-\phi)M(\phi)} |(\mathcal{K}(t, \psi(t)) - \mathcal{K}(t, \tilde{\psi}(t)))| \\
329 \quad & \quad + \frac{2\phi}{(2-\phi)M(\phi)} \int_0^t |(\mathcal{K}(\tau, \psi(\tau)) - \mathcal{K}(\tau, \tilde{\psi}(\tau)))| d\tau
\end{aligned}$$

330 Using Lipschitz's inequality (33), we get the following:

$$\begin{aligned}
331 \quad & |\psi(t) - \tilde{\psi}(t)| \\
332 \quad & \leq \left| \psi(t) - \tilde{\psi}_0 - \frac{2(1-\phi)}{(2-\phi)M(\phi)} \mathcal{K}(t, \psi(t)) - \frac{2\phi}{(2-\phi)M(\phi)} \int_0^t \mathcal{K}(\tau, \psi(\tau)) d\tau \right| \\
333 \quad & \quad + \frac{2L(1-\phi)}{(2-\phi)M(\phi)} |\psi(t) - \tilde{\psi}(t)| + \frac{2L\phi}{(2-\phi)M(\phi)} \int_0^t |\psi(\tau) - \tilde{\psi}(\tau)| d\tau \\
334 \quad & = \left| \psi(t) - \tilde{\psi}_0 - \frac{2(1-\phi)}{(2-\phi)M(\phi)} \mathcal{K}(t, \psi(t)) - \frac{2\phi}{(2-\phi)M(\phi)} \int_0^t \mathcal{K}(\tau, \psi(\tau)) d\tau \right| \\
335 \quad & \quad + \frac{2L((1-\phi)+\phi T_{\text{sup}})}{(2-\phi)M(\phi)} |\psi(t) - \tilde{\psi}(t)|
\end{aligned}$$

336 Using (31) and (37), we get,

$$\begin{aligned}
337 \quad & |\psi(t) - \tilde{\psi}(t)| \leq \frac{\eta \mathcal{Y}(t)}{(2-\phi)M(\phi)} + \theta |\psi(t) - \tilde{\psi}(t)| \\
338 \quad & \leq \frac{\eta \mathcal{Y}(t)}{(1-\theta)(2-\phi)M(\phi)} = \varepsilon_\phi \mathcal{Y}(t)
\end{aligned}$$

339

340 7. Numerical Scheme

341 In this section, a new numerical scheme is obtained to solve numerically the
342 fractional-order system representing the proposed model. Consider the fractional-order
343 equation ${}^C D_t^\phi \psi(t) = \mathcal{K}(t, \psi(t))$, applying the fundamental theorem of fractional calculus
344 yields; an iterative scheme is obtained as follows:

$$345 \quad \psi(t_{n+1}) - \psi(0) = \frac{2(1-\phi)}{(2-\phi)M(\phi)} \mathcal{K}(t_n, \psi(t_n)) + \frac{2\phi}{(2-\phi)M(\phi)} \int_0^{t_{n+1}} \mathcal{K}(\tau, \psi(\tau)) d\tau \quad (38)$$

346 Replacing value of $\psi(t_n)$, we get,

$$\begin{aligned}
347 \quad \psi(t_{n+1}) &= \psi(t_n) + \frac{2(1-\phi)}{(2-\phi)M(\phi)} [\mathcal{K}(t_n, \psi(t_n)) - \mathcal{K}(t_{n-1}, \psi(t_{n-1}))] \\
348 \quad &+ \frac{2\phi}{(2-\phi)M(\phi)} \int_{t_n}^{t_{n+1}} \mathcal{K}(\tau, \psi(\tau)) d\tau \quad (39)
\end{aligned}$$

349 Considering uniform step-size h along **the** time axis, the integral can be approximated as in
350 **the** classical two-step Adams-Bashforth scheme as follows:

$$351 \quad \int_{t_n}^{t_{n+1}} \mathcal{K}(\tau, \psi(\tau)) d\tau \approx \frac{3h}{2} \mathcal{K}(t_n, \psi(t_n)) - \frac{h}{2} \mathcal{K}(t_{n-1}, \psi(t_{n-1})) \quad (40)$$

352 Substituting **the** value of approximated integral (40) into **the** above equation (39),

$$\begin{aligned}
353 \quad \psi(t_{n+1}) &= \psi(t_n) + \frac{2(1-\phi)}{(2-\phi)M(\phi)} [\mathcal{K}(t_n, \psi(t_n)) - \mathcal{K}(t_{n-1}, \psi(t_{n-1}))] \\
354 \quad &+ \frac{2\phi}{(2-\phi)M(\phi)} \left[\frac{3h}{2} \mathcal{K}(t_n, \psi(t_n)) - \frac{h}{2} \mathcal{K}(t_{n-1}, \psi(t_{n-1})) \right]
\end{aligned}$$

355 Since, $M(\phi)$ is a normalizing function with $M(0) = M(1) = 1$. So, let us assume $M(\phi) =$
356 $(2 - \phi^2)/(2 - \phi)$ which satisfies $M(0) = M(1) = 1$. Thus,

$$357 \quad \psi(t_{n+1}) = \psi(t_n) + \frac{2+(3h-2)\phi}{2-\phi^2} \mathcal{K}(t_n, \psi(t_n)) - \frac{2+(h-2)\phi}{2-\phi^2} \mathcal{K}(t_{n-1}, \psi(t_{n-1})) \quad (41)$$

358 Hence, the fractional-order model (3) has the following numerical scheme to obtain the
359 numerical solutions.

$$360 \quad \vec{\psi}(t_{n+1}) = \vec{\psi}(t_n) + \frac{2+(3h-2)\phi}{2-\phi^2} \vec{\mathcal{K}}(t_n, \vec{\psi}(t_n)) - \frac{2+(h-2)\phi}{2-\phi^2} \vec{\mathcal{K}}(t_{n-1}, \vec{\psi}(t_{n-1})) \quad (42)$$

361

362 7.1. Numerical Simulations

363 This section uses MATLAB software to perform the numerical simulations from the
364 obtained numerical scheme (42). The total initial population is assumed to be $N(0) = 100$
365 and the initial values of the compartments are assumed to be $S(0) = 80, E(0) = 10, I(0) =$
366 $5, H(0) = 3, D(0) = 2, R(0) = 0$. The used values of the parameters are as follows: $B = 10,$
367 $\mu = 0.1, \alpha_1 = 0.75, \alpha_2 = 0.85, \alpha_3 = 0.425, \alpha_4 = 0.2, \alpha_5 = 0.15,$ and $\alpha_6 = 0.25$.

368 Figure 2 shows the transmission dynamics of each compartment listed in the model
369 over time. The behavior is smooth, and it validates the theoretical results. Figure 3 shows the
370 behavior for different values of fractional order. For relatively small values of the fractional
371 order, the number of infectious individuals reaches the exact peak of approximately 19 cases
372 but takes a relatively long time.

373 Figure 4 shows the behavior of the Ebola infectious cases for the different values of
374 the contact rate over time. The relatively more contact rate of susceptible with the pathogen
375 carriers individuals will surge the number of Ebola infectious individuals. The contact rate is
376 the crucial parameter for this model that directly influences the cases of Ebola. It shows that
377 the most efficient way to control the spread of EBOV infection is to control the contact rate
378 parameter.

379 Figure 5 and Figure 6 illustrate the behavior of the Susceptible $S(t)$ and Deceased
380 $D(t)$ for various values of contact rate α_1 over time, respectively.

381 In Table 2, the CPU time usage is listed with the different values of step size Δt and
382 iterations n of the mentioned numerical scheme for this proposed model. The table makes it
383 clear that the proposed strategy increases efficiency while taking less time.

384

385 8. Conclusions

386 In this study, an epidemic model for the Ebola disease is formulated using the Caputo-
387 Fabrizio fractional derivative. The basic reproduction number (\mathcal{R}_0) is calculated using the
388 next-generation matrix approach. It analyzed the condition for the existence and uniqueness
389 of the model's solution using the Fixed-point theorem approach. Additionally, the stability
390 condition for the suggested model's generalized Ulam-Hyers-Rassias stability is found. It

391 illustrates how the approximate solution of the proposed model differs for integer and
392 fractional orders in numerical simulations. Additionally, it displays the behavior of Ebola
393 infections in deceased and vulnerable individuals at various contact rate values. In future, the
394 authors can study this approach for other infectious diseases to get better insight about the
395 transmission of the diseases whose outcomes may help medical fraternity to work effectively.

396

397 **9. Declaration of Competing Interest**

398 The authors declare that they have no known competing financial interests or personal
399 relationships that could have appeared to influence the work reported in this paper.

400

401 **10. Credit authorship contribution statement**

402 **Shah Nita:** gave the concept, supervision. **Chaudhary Kapil:** wrote the manuscript,
403 methodology, analyzed model mathematically and software.

404

405 **11. Acknowledgment**

406 The authors thank Editor of the journal and blind reviewers for their constructive
407 comments on the research. Second author, Kapil Chaudhary is supported by CSIR-
408 JRF(Council of Scientific and Industrial Research- Junior research fellowship) File No.
409 09/0070(13467)/2022-EMR-I.

410

411 **References**

412 Bisimwa, P., Biamba, C., Aborode, A. T., Cakwira, H., & Akilimali, A. (2022). Ebola virus
413 disease outbreak in the Democratic Republic of the Congo: A mini-review. *Annals*

414 *of Medicine and Surgery*, 80, 104213.

415 Diekmann, O., Heesterbeek, J. A. P., & Metz, J. A. (1990). On the definition and the
416 computation of the basic reproduction ratio R_0 in models for infectious diseases in
417 heterogeneous populations. *Journal of mathematical biology*, 28(4), 365-382.

418 Feldmann, H., Sprecher, A., & Geisbert, T. W. (2020). Ebola. *New England Journal of*
419 *Medicine*, 382(19), 1832-1842.

420 Gao, F., Li, X., Li, W., & Zhou, X. (2021). Stability analysis of a fractional-order novel
421 hepatitis B virus model with immune delay based on Caputo-Fabrizio
422 derivative. *Chaos, Solitons & Fractals*, 142, 110436.

423 Hammouch, Z., Rasul, R. R., Ouakka, A., & Elazzouzi, A. (2022). Mathematical analysis
424 and numerical simulation of the Ebola epidemic disease in the sense of
425 conformable derivative. *Chaos, Solitons & Fractals*, 158, 112006.

426 Hussain, A., Baleanu, D., & Adeel, M. (2020). Existence of solution and stability for the
427 fractional order novel coronavirus (nCoV-2019) model. *Advances in Difference*
428 *Equations*, 2020(1), 1-9.

429 Khajehsaeid, H. (2018). A comparison between fractional-order and integer-order
430 differential finite deformation viscoelastic models: Effects of filler content and
431 loading rate on material parameters. *International Journal of Applied Mechanics*,
432 10(09), 1850099.

433 Liu, K., Fečkan, M., O'Regan, D., & Wang, J. (2019). Hyers–Ulam stability and existence
434 of solutions for differential equations with Caputo–Fabrizio fractional
435 derivative. *Mathematics*, 7(4), 333.

436 Liu, K., Fečkan, M., & Wang, J. (2020). A Fixed-Point Approach to the Hyers–Ulam

437 Stability of Caputo–Fabrizio Fractional Differential Equations. *Mathematics*, 8(4),
438 647.

439 Losada, J., & Nieto, J. J. (2015). Properties of a new fractional derivative without singular
440 kernel. *Progress in Fractional Differentiation and Application*, 1(2), 87-92.

441 Nwajeri, U. K., Omame, A., & Onyenegecha, C. P. (2021). Analysis of a fractional order
442 model for HPV and CT co-infection. *Results in Physics*, 28, 104643.

443 Nwajeri, U. K., Panle, A. B., Omame, A., Obi, M. C., & Onyenegecha, C. P. (2022). On the
444 fractional order model for HPV and Syphilis using non–singular kernel. *Results in*
445 *Physics*, 37, 105463.

446 Otunuga, O. M. (2021). Estimation of epidemiological parameters for COVID-19 cases
447 using a stochastic SEIRS epidemic model with vital dynamics. *Results in*
448 *Physics*, 28, 104664.

449 Shah, N. H., Patel, Z. A., & Yeolekar, B. M. (2019). Vertical dynamics of Ebola with
450 media impact. *Journal of King Saud University-Science*, 31(4), 567-574.

451 Shaikh, A. S., & Nisar, K. S. (2019). Transmission dynamics of fractional order Typhoid
452 fever model using Caputo–Fabrizio operator. *Chaos, Solitons & Fractals*, 128,
453 355-365.

454 Singh, H. (2020). Analysis for fractional dynamics of Ebola virus model. *Chaos, Solitons &*
455 *Fractals*, 138, 109992.

456 Singh, H., Baleanu, D., Singh, J., & Dutta, H. (2021). Computational study of fractional
457 order smoking model. *Chaos, Solitons & Fractals*, 142, 110440.

458 Singh, H., Srivastava, H. M., Hammouch, Z., & Nisar, K. S. (2021). Numerical simulation
459 and stability analysis for the fractional-order dynamics of COVID-19. *Results in*

460 *physics*, 20, 103722.

461 Solís-Pérez, J. E., Gómez-Aguilar, J. F., & Atangana, A. (2018). Novel numerical method
462 for solving variable-order fractional differential equations with power, exponential
463 and Mittag-Leffler laws. *Chaos, Solitons & Fractals*, 114, 175-185.

464 Srivastava, H. M., & Saad, K. M. (2020). Numerical simulation of the fractal-fractional
465 Ebola virus. *Fractal and Fractional*, 4(4), 49.

466 World Health Organization. (2014). *What we know about transmission of the ebola virus*
467 *among humans*. World Health Organization. Retrieved August 4, 2022, from
468 <http://www.who.int/mediacentre/news/ebola/06-october-2014/en/>

469 Zhang, Z., & Jain, S. (2020). Mathematical model of Ebola and Covid-19 with fractional
470 differential operators: Non-Markovian process and class for virus pathogen in the
471 environment. *Chaos, Solitons & Fractals*, 140, 110175.

472




Figure 1 The flow diagram of the proposed model.




Figure 2 Smooth behavior of compartments over time using integer-order $\phi = 1$.




Figure 3 Effect of different values of fractional order ϕ on infectious individuals $I(t)$ over time t .




Figure 4 Effect of the contact rate α_1 on infectious individuals $I(t)$ over time t with integer-order $\phi = 1$.




Figure 5 Effect of the contact rate α_1 on susceptible individuals $S(t)$ over time t with integer-order $\phi = 1$.




Figure 6 Effect of the contact rate α_1 on deceased individuals $D(t)$ over time t with integer order $\phi = 1$.

Parameter	Meaning
B	The birth rate of the population
α_1	Contact rate of susceptible with EBOV carriers
α_2	The transmission rate of exposed getting infectious from the Ebola
α_3	The transmission rate of infectious getting hospitalized
α_4	The transmission rate of infectious getting critically ill or deceased
α_5	The transmission rate of hospitalized getting critically ill or deceased
α_6	The transmission rate of hospitalized getting recovered
μ	The death rate of the population

Table 1 Meaning of parameters used in the proposed model.

Step size (Δt)	Number of iterations (n)	CPU time (s)
0.1	100	0.29
0.01	1000	0.34
0.001	10^4	0.41
0.0001	10^5	1.31
0.00001	10^6	2.39

Table 2 CPU time usage of the different values of Δt and n .