

Keywords Index

A

ab initio; 37 (3), 327-343
absolute growth rate; 37 (1), 21-27
absorption; 37 (1), 65-72
Acinetobacter; 37 (1), 7-13
activated sludge; 37 (1), 55-63
acute toxicity; 37 (4), 433-439
age structure; 37 (4), 485-491
Aglaia andamanica; 37 (1), 37-41
air traffic management; 37 (3), 345-351
airplane allocation; 37 (3), 345-351
airplane routing; 37 (3), 345-351
Allotropa suasardi Sarkar & Polaszek; 37 (4), 381-387
amino acids; 37 (4), 397-400
animal model; 37 (1), 21-27
antibacterial; 37 (1), 43-48; 37 (3), 309-317
antibiotic-resistant enterococci; 37 (3), 299-307
anticancer; 37 (3), 309-317
anti-carcinogenic; 37 (3), 275-282
anti-inflammation; 37 (1), 43-48
antimicrobial activity; 37 (5), 587-594
antimicrobial; 37 (3), 275-282
antinutritional compounds; 37 (5), 507-514
antioxidant effect; 37 (2), 163-169
application methods; 37 (6), 635-642
arrowroot starch; 37 (2), 193-199
asperity; 37 (6), 683-690
associative memory; 37 (6), 669-674
atretic follicles; 37 (5), 569-573
autonomous underwater robot; 37 (6), 659-668

B

B. juncea growth; 37 (6), 631-634
bacterial cells; 37 (6), 625-630
bacterial community; 37 (3), 291-297
bacterial determinants; 37 (1), 29-36
Bactrocera cacuminata; 37 (6), 617-624
Bactrocera tryoni; 37 (6), 617-624
bee algorithm; 37 (2), 221-230
behavior; 37 (4), 397-400
beta exponentiated weibull poisson; 37 (1), 97-108
beta-g distribution; 37 (1), 97-108

betaine aldehyde dehydrogenase2; 37 (5), 533-537
bifurcation; 37 (6), 719-725
biodegradable polymer; 37 (2), 193-199
biodiesel; 37 (4), 389-395
bioemulsifier; 37 (4), 425-432
biogas; 37 (1), 65-72
biological; 37 (3), 327-343
bioreactor; 37 (1), 73-80
block diagonal structure; 37 (4), 477-484
Boro rice; 37 (4), 455-464
bryophyte; 37 (4), 409-415
buffalo leech; 37 (5), 581-585
buffers; 37 (6), 625-630

C

callus culture; 37 (1), 49-53
carbazole alkaloid; 37 (1), 43-48
carbon dioxide; 37 (1), 65-72
cetane number; 37 (4), 389-395
chicken; 37 (3), 299-307
cholesterol; 37 (3), 275-282
cisplatin; 37 (2), 155-162
Clausena lansium; 37 (1), 43-48
coastal area; 37 (2), 147-153
Coastal Bangladesh; 37 (4), 455-464
Cocos nucifera L.; 37 (4), 417-423
Coix lacryma-jobi L.; 37 (4), 441-447
composition; 37 (2), 147-153
conceptual design; 37 (6), 659-668
concrete; 37 (2), 201-207
Corbiculidae; 37 (1), 15-20
corn downy mildew; 37 (5), 515-522
corn; 37 (5), 515-522
correlation analysis; 36 (4), 209-219
cotton dust ash; 37 (2), 201-207
count data; 37 (6), 701-711
crab; 37 (2), 135-145
Croton stellatopilosus; 37 (1), 49-53
crunch training; 37 (5), 545-551
crustacean; 37 (2), 135-145
crystallization; 37 (3), 353-359
culled dairy cow; 37 (2), 109-117

culture system; 37 (5), 575-580
 cyclic loading; 37 (6), 683-690
Cyclosoorus terminans; 37 (3), 309-317
 ε -chainable; 37 (1), 89-92
 ε -contractive; 37 (1), 89-92

D

daily precipitation; 37 (1), 81-88
 daily rainfall; 37 (1), 81-88
 degradation; 37 (1), 65-72
 DEM; 37 (6), 691-700
 demographic transition; 37 (4), 485-491
 denitrification; 37 (1), 73-80
 density; 37 (5), 581-585
 diet; 37 (2), 109-117
 digestive tract; 37 (5), 561-567
 dilation; 37 (6), 683-690
 di-nucleotide SSR; 37 (2), 119-127
 dolomite; 37 (6), 643-650
 dry rubber content; 37 (6), 713-718
 duckweed; 37 (5), 575-580

E

ecorace; 37 (5), 499-505
 electrical resistivity tomography (ERT); 37 (5), 675-681
 elicitation; 37 (1), 49-53
 encapsulation; 37 (4), 425-432
 endogenous hormonal status; 37 (5), 539-544
 endothelial function; 37 (5), 545-551
 energy input-output; 37 (4), 455-464
Enterobacter asburiae strain RS83; 37 (1), 29-36
 enterocin; 37 (3), 299-307
Enterococcus faecium; 37 (3), 299-307
 eri silkworm; 37 (4), 401-408; 37 (5), 499-505
 ethanol; 37 (5), 533-537
 ethnobotany; 37 (3), 369-379
 excessive zero; 37 (6), 701-711

F

factor analysis; 37 (1), 81-88; 37 (4), 485-491
 farm categories; 37 (4), 455-464
 fatty acid and carotenoid levels; 37 (6), 609-616
 fatty acid composition; 37 (4), 389-395
 fatty acid methyl ester; 37 (4), 389-395
 fatty acid; 37 (2), 109-117; 37 (5), 507-514
 feeding frequency; 37 (2), 129-134
 female mud crab; 37 (6), 609-616
 fertilizer; 37 (6), 643-650
 fingerprint image; 37 (2), 241-246
 fire rating; 37 (2), 183-192
 fish; 37 (1), 1-5; 37 (5), 569-573
 flood map; 37 (2), 171-181
 flowering time; 37 (3), 247-254
 flow-mediated dilatation; 37 (5), 545-551
 fracture stiffness; 37 (6), 683-690
 fruit juices; 37 (3), 265-270

fruiting body; 37 (4), 433-439
 fumigation; 37 (5), 523-531
 fusion; 37 (6), 719-725
 Fuzzy 3- metric space; 37 (1), 89-92
 fuzzy bitopological spaces; 37 (1), 93-96
 fuzzy *bLO* sets; 37 (1), 93-96

G

gallic acid; 37 (3), 353-359
 gamma regression; 37 (1), 81-88
 generalized confidence interval; 37 (2), 231-240
 genetic algorithm; 37 (2), 221-230
 genetic parameters; 37 (1), 21-27
 geranylgeraniol; 37 (1), 49-53
 germination index; 37 (5), 553-559
 ginseng extract G115; 37 (2), 163-169
 GIS; 37 (4), 493-498; 37 (6), 691-700
 GLM; 37 (1), 81-88
 global warming; 37 (3), 369-379
 glucosinolate; 37 (6), 625-630
 growth; 37 (5), 581-585
 gut bacteria community; 37 (6), 617-624

H

habitat selection; 37 (3), 255-263
Haliotis mariae; 37 (3), 255-263
 Hamacher T-co-norm; 37 (2), 241-246
 heat affected zone; 37 (3), 361-367
 heat input; 37 (3), 361-367
 heavy metal; 37 (6), 651-657
Hemibagrus filamentus; 37 (1), 1-5
 high temperature; 37 (4), 401-408
 high-dimensional data; 37 (4), 477-484
 hirudinea manillensis; 37 (5), 581-585
 histochemistry; 37 (5), 561-567
 histology; 37 (5), 561-567; 37 (5), 569-573
 histopathology; 37 (1), 1-5
 hornworts; 37 (3), 271-274
 host preference; 37 (4), 381-387
 host stage selection; 37 (4), 381-387
 host suitability; 37 (4), 381-387
 house of quality; 37 (6), 659-668
 human skin; 37 (1), 7-13
 hypothesis testing; 37 (4), 477-484
 hypouricemia; 37 (4), 441-447
 hypoxanthine; 37 (3), 283-289

I

500°C isotherm method; 37 (2), 183-192
 image enhancement; 37 (2), 241-246
 inbred line; 37 (5), 515-522
 indole acetic acid; 37 (5), 553-559
 industrial waste; 37 (2), 201-207
 inhibitor; 37 (3), 327-343
 injection molding; 37 (2), 193-199
 inorganic phosphorus; 37 (5), 515-522

insecticidal; 37 (5), 523-531
 integer programming; 37 (3), 345-351
 interruptins A-C; 37 (3), 309-317
 intuitionistic fuzzy (soft) ideals; 37 (4), 465-475
 intuitionistic fuzzy (soft) subalgebras; 37 (4), 465-475
 intuitionistic fuzzy set; 37 (2), 241-246
 iron filings; 37 (1), 55-63
 isothiocyanate; 37 (6), 625-630
 ISSR marker; 37 (5), 499-505

J

jackfruit seeds; 37 (3), 353-359

K

kefir; 37 (3), 275-282
 kidney; 37 (1), 1-5
Kung-Som; 37 (3), 291-297

L

Lactobacillus paracasei SD1; 37 (3), 265-270
Lactobacillus plantarum; 37 (4), 425-432
 land use land cover; 37 (6), 691-700
 landsat; 37 (6), 691-700
 large search space; 37 (2), 221-230
 larviculture; 37 (2), 135-145
 laser welding; 37 (3), 361-367
 lateritic soil; 37 (1), 55-63
 latex; 37 (6), 713-718
 leg raise training; 37 (5), 545-551
Lentinus polychrous; 37 (4), 433-439
L-H transition; 37 (6), 719-725
 light integral; 37 (3), 247-254
 linkage map; 37 (2), 119-127
 lipase; 37 (1), 7-13
 lipid oxidation; 37 (2), 109-117
 liquid endosperm; 37 (5), 533-537
 lithium chloride; 37 (5), 533-537
 liver; 37 (1), 1-5
 liverworts; 37 (4), 409-415
 low humidity; 37 (4), 401-408

M

macrophage migration inhibitory factor (MIF);
 37 (4), 417-423
 Malaysia; 37 (2), 147-153
Mangifera; 37 (2), 119-127
 mangrove clam; 37 (1), 15-20
 marble goby; 37 (4), 397-400
 mass culture; 37 (5), 575-580
 mathematical model; 37 (5), 595-600
 maximum likelihood estimation; 37 (6), 701-711
 mean vector; 37 (4), 477-484
 Meliaceae; 37 (1), 37-41
 menaquinone-7; 37 (3), 283-289
 method of moments; 37 (6), 701-711
 methyl bromide; 37 (5), 523-531

methyl oleate synthesis; 37 (1), 7-13
 microencapsulation; 37 (3), 265-270
 microwave; 37 (6), 713-718
 molecular cloning; 37 (6), 617-624
 monoethanolamine; 37 (1), 65-72
 Monte Carlo simulation; 37 (2), 231-240
 moringa oleifera seed; 37 (5), 587-594
 morphometric analysis; 37 (1), 15-20
 motorway; 36 (4), 209-219
 multiple regression; 36 (4), 209-219
 mungbean starch; 37 (2), 193-199
 municipal solid waste; 37 (4), 449-454
 municipality; 37 (4), 449-454
 Murashige and Skoog 2; 37 (5), 553-559
 mushroom; 37 (4), 433-439
 mycelia; 37 (4), 433-439

N

1-naphthol; 37 (3), 283-289
 Na^+/K^+ ratio; 37 (6), 631-634
 nasturtium; 37 (3), 247-254
 NBO; 37 (3), 327-343
 Nepal; 37 (4), 485-491
 nephroprotection; 37 (4), 441-447
 nephroprotective; 37 (2), 155-162
 nitrification; 37 (1), 73-80
 nitrile; 37 (6), 625-630
 nitrogen; 37 (1), 73-80
 NMR; 37 (3), 327-343
 non-invasive geophysical methods; 37 (5), 675-681
Notothylas yunannensis; 37 (3), 271-274
 nucleic acid; 37 (5), 533-537

O

occasionally weakly compatible; 37 (1), 89-92
 oil; 37 (5), 587-594
 one step purification; 37 (1), 7-13
 organoleptic properties; 37 (2), 109-117
 ovariectomy; 37 (4), 417-423
 ovary maturation; 37 (6), 609-616
 overdispersion; 37 (6), 701-711
 oxidative stability; 37 (5), 587-594
Oxyeleotris marmoratus; 37 (4), 397-400

P

P- fertilizer; 37 (6), 631-634
 parkinsonism; 37 (2), 163-169
 pattern recognition; 37 (6), 669-674
 PCR-DGGE; 37 (3), 291-297
 periodontal disease; 37 (1), 43-48
 permeable reactive barrier; 37 (1), 55-63
Phenacoccus manihoti Matile-Ferrero; 37 (4), 381-387
 phosphate solubilizing bacteria; 37 (6), 651-657
 photoperiod; 37 (3), 247-254
 Phyllanthin; 37 (3), 319-326
Phyllanthus amarus; 37 (3), 319-326

phytic acid; 37 (5), 515-522
 phytotoxicity; 37 (5), 523-531
 plant growth enhancement; 37 (1), 29-36
 plasma; 37 (6), 719-725
 porous combustion; 37 (5), 595-600
Portunus pelagicus; 37 (2), 129-134; 37 (2), 135-145
 potassium oxonate; 37 (4), 441-447
 prebiotics; 37 (3), 353-359
 precursor addition; 37 (1), 49-53
 pre-harvest fruit drop; 37 (5), 539-544
 premixed combustion; 37 (5), 595-600
 prime fuzzy soft bi-ideal; 37 (5), 601-608
 probiotic; 37 (3), 265-270; 37 (3), 275-282; 37 (4), 425-432
 protein pattern; 37 (5), 507-514
 pummelo; 37 (5), 539-544

Q

QFD; 37 (6), 659-668

R

raster model; 37 (2), 171-181
Rastrelliger brachysoma; 37 (5), 561-567; 37 (5), 569-573
 rating curve; 37 (2), 171-181
 RAW264.7 cells; 37 (1), 37-41
 RBL-2H3 cells; 37 (1), 37-41
 ready mixed concrete truck scheduling; 37 (2), 221-230
 rearing techniques; 37 (2), 135-145
 reinforced concrete; 37 (2), 183-192
 relative growth rate; 37 (1), 21-27
 remote sensing; 37 (6), 691-700
 renal epithelial cell; 37 (2), 155-162
 reproduction; 37 (5), 581-585
 residual stress; 37 (3), 361-367
 restocking; 37 (3), 255-263
 rhizosphere; 37 (5), 553-559
 River Ganga; 37 (6), 651-657
 road users' satisfaction level; 36 (4), 209-219
 roasting; 37 (5), 507-514
 robot design; 37 (6), 659-668
 ROS scavenging; 37 (3), 309-317
 ROS; 37 (2), 155-162
 rotenone; 37 (2), 163-169
rpoB gene; 37 (3), 291-297
 rubber plantation; 37 (6), 643-650
 rural livelihoods; 37 (3), 369-379

S

16SrRNA gene; 37 (6), 617-624
 salinity classes; 37 (4), 455-464
 salt stress; 37 (6), 631-634
 Sarawak; 37 (1), 15-20; 37 (2), 147-153
Schistochila; 37 (4), 409-415
 Schistochilaceae; 37 (4), 409-415
 screening; 37 (5), 499-505
 seaweeds; 37 (2), 147-153
 seed densities; 37 (3), 255-263

seeding; 37 (3), 255-263
 seepage; 37 (5), 675-681
 selection; 37 (4), 401-408
 semiprime fuzzy soft bi-ideal; 37 (5), 601-608
 sex structure; 37 (4), 485-491
 shade; 37 (3), 247-254
 shear strength; 37 (6), 683-690
 sheep; 37 (1), 21-27
 shelter; 37 (2), 129-134
 shield metal arc welding; 37 (3), 361-367
 short mackerel; 37 (5), 569-573
 side crunch training; 37 (5), 545-551
 silymarin; 37 (2), 155-162
 simple confidence interval for lognormal means;
 37 (2), 231-240
 simple confidence interval for ratio of lognormal means;
 37 (2), 231-240
 soft sets; 37 (4), 493-498
 soft spatial region; 37 (4), 493-498
 soft topology; 37 (4), 493-498
 soil erosion; 37 (6), 691-700
 soil; 37 (6), 635-642
 solid phase extraction; 37 (3), 353-359
 species distribution modeling; 37 (3), 369-379
 spent yeast; 37 (4), 425-432
 spore; 37 (3), 271-274
 stem cell proliferation; 37 (3), 309-317
 strongly prime fuzzy soft bi-ideal; 37 (5), 601-608
 structural elements; 37 (2), 183-192
 Subtraction algebras; 37 (4), 465-475
 survival rate of crab larvae; 37 (2), 129-134
 survivorship; 37 (5), 581-585

T

tabulated data; 37 (2), 183-192
 Tapee River; 37 (1), 1-5
 tapping rubber; 37 (6), 643-650
 taste preferences; 37 (4), 397-400
 taxonomy; 37 (3), 271-274
 temperature; 37 (3), 247-254
 Tenera palm oil; 37 (4), 389-395
 Thai Navy; 37 (6), 659-668
 Thailand; 37 (3), 271-274; 37 (4), 409-415; 37 (4), 449-454;
 37 (5), 561-567; 37 (5), 569-573
 The Center-TRACON Automation System; 37 (3), 345-351
 thermoplastic starch; 37 (2), 193-199
 Thong Dee; 37 (5), 539-544
 TLC-densitometry; 37 (3), 319-326
 TLC-image analysis; 37 (3), 319-326
 tokamak; 37 (6), 719-725
 tolerance; 37 (4), 401-408
 tomato seedlings; 37 (5), 553-559
 transient thermal analysis; 37 (2), 183-192
 transmission electron microscopy (TEM); 37 (4), 417-423
 trigonometric chaotic neuron; 37 (6), 669-674
Tropaeolum majus L.; 37 (3), 247-254

tropical forest; 37 (3), 369-379
tween 80; 37 (3), 283-289
type II fuzzy set; 37 (2), 241-246

U

ultrasonication; 37 (3), 283-289
uncertain factors; 37 (2), 221-230
unknown bridge foundation determination; 37 (5), 675-681
Upper Ping River; 37 (2), 171-181

V

V3 region; 37 (3), 291-297
Validation; 37 (3), 319-326
vincristine; 37 (3), 327-343
vorticity-stream function; 37 (5), 595-600

W

waste collection costs; 37 (4), 449-454
waste generation; 37 (4), 449-454
wastewater; 37 (1), 73-80

water meal; 37 (5), 575-580
watershed; 37 (6), 691-700
waveguide; 37 (6), 713-718
weakly commuting; 37 (1), 89-92
wetland; 37 (1), 15-20
wheat; 37 (6), 635-642
Wolffia globosa; 37 (5), 575-580

X

xanthine oxidase; 37 (4), 441-447

Y

yield; 37 (5), 499-505
young-coconut juice; 37 (4), 417-423

Z

zero inflated negative binomial-Crack distribution;
37 (6), 701-711
zinc; 37 (1), 55-63