

*Original Article*

# A numerical approach to the solution of nonlinear Volterra-Fredholm integro-differential equations

Ganiyu Ajileye<sup>1\*</sup>, Ojo Olamiposi Aduroja<sup>2</sup>, Ineye Grace Amakoromo<sup>1</sup>,  
and Sikiru Adesina Amoo<sup>1</sup>

<sup>1</sup> *Department of Mathematics and Statistics, Federal University Wukari,  
Wukari, Taraba, 670102 Nigeria*

<sup>2</sup> *Department of Mathematics, University of Ilesa, Ilesa, Osun, Nigeria*

Received: 27 November 2023; Revised: 14 December 2023; Accepted: 27 December 2023

---

**Abstract**

In this study, a numerical method is presented for solving first-order Volterra-Fredholm Integro-differential Equations (V-FIDEs). The approach converts V-FIDEs into integral equations and uses power series polynomials to approximate them. The modeled problem is transformed into a system of algebraic equations, and then it is solved using the standard collocation method. Numerical examples are utilized to assess the method's effectiveness after the approach's uniqueness and convergence have been established. The results demonstrate that the method competes favorably with other methods.

**Keywords:** collocation method, Volterra-Fredholm, integro-differential equations, approximate solution, polynomial power series

---

**1. Introduction**

Integro-differential Equations (IDEs) are strong tools in pure and applied mathematics, engineering, and physics. IDEs are used in many mathematical representations of physical phenomena in fluid dynamics, heat transfer, diffusion processes, neutron diffusion, biological models, nano-hydrodynamics, economics, and population growth models. In 1926, Vito Volterra used an integro-differential approach to explore population increase with a focus on hereditary effects (Rahman, 2007).

The various methods for finding the numerical solution of IDEs include Bernstein method (Irfan, Kumar, & Kapoor, 2014), Adomian decompositions method (Khan & Bakoda, 2013; Mittal & Nigam, 2008), Finite difference-Simpson method (Garba & Bichi, 2021), collocation method (Agbolade & Anake, 2017; Ajileye & Aminu, 2022; Ajileye &

Amoo, 2023; Ajileye, James, Ayinde & Oyedepo 2022; Nemati, Lima & Ordok-hani, 2015), the Modified Adomian Decomposition Method (MADM) (Ahmad, Ullah, Ullah, Ahmad, Shah & Ahmad, 2021), hybrid linear multistep method (Mehdiyera, Ibrahim & Imanova, 2019; Mehdiyera, Imanova & Ibrahim, 2015), Chebyshev-Galerkin method (Issa & Saleh, 2017), Bernoulli matrix method (Bhraway, Tohidi, & Soleymani, 2012), differential transform method (Ercan & Kharerah, 2013), Haar collocation method (Amin, Shah, Mlaiki, Yuzbasi, Abdeljawad, & Hussain, 2022), Lagrange interpolation (Shahsavaran & Shahsavaran, 2012), differential transformation (Darania & Ebadian, 2007), block pulse functions' operational matrices (Rahmani, Rahimin, & Mordad, 2011), Chebyshev polynomials (Maadadi & Rahmoune, 2018), Optimal Auxiliary Function Method (OAFM) (Zada, Al-Hamami, Nawaz, Jehanzeb, Morsy, Abdei-Aty, & Nisar, 2021), and spectral homotopy method (Atabakan, Nasab, Kilicman, & Eshkuvatov, 2013). Hou, Niu, Xu & Ngolo (2021) presented a method combining the simplified reproducing kernel method and the homotopy perturbation method to solve nonlinear Volterra-Fredholm integro-differential equations. They converted nonlinear to linear using

---

\*Corresponding author

Email address: [ajileye@fuwukari.edu.ng](mailto:ajileye@fuwukari.edu.ng)

the homotopy perturbation method and adopted the simplified reproducing kernel method to solve the linear problem.

In this paper, we present a collocation method for the numerical solution of first-order V-FIDEs of the form:

$$y'(x) + q(x)y(x) + \lambda_1 \int_0^x k_1(x,t) F_1(y(t))dt + \lambda_2 \int_0^1 k_2(x,t) F_2(y(t))dt = g(x) \tag{1}$$

with the initial condition,

$$y(a) = c \tag{2}$$

where  $k_1(x, t)$  and  $k_2(x, t)$  are the Volterra and Fredholm integral kernel functions, respectively, and  $\lambda^1, \lambda^2, a, c,$  and  $q$  are known constants.  $g(x)$  is a given function, and  $y(x)$  is the unknown function to be determined.

### 2. Basic Definitions and Terms

We give certain definitions and fundamental notions in this section for the purpose of problem formulation.

**Definition 1.** (Ajileye & Amoo, 2023) Let  $(a_m), m \geq 0$  be a sequence of real numbers. The power series  $w$  with coefficients  $a_m$  is an expression.

$$y(w) = \sum_{m=0}^M a_m w^m = \Phi(w)A \tag{3}$$

where

$$\Phi(w) = [1 \ w \ w^2 \ \dots \ w^M], \ A = [a_0 \ a_1 \ \dots \ a_M]^T$$

**Definition 2.** (Agbolade & Anake, 2017) The desired collocation points within an interval are determined using this method. For interval  $[a, b]$  they are provided by

$$l_u = a + \frac{(b-a)u}{M}, \ u = 0, 1, 2 \dots \dots M \tag{4}$$

**Definition 3.** (Ajileye, James, Ayinde, & Oyedepo, 2022) Let  $z(s)$  be an integrable function, then

$${}_0I_x^\beta(z(s)) = \frac{1}{\Gamma(\alpha)} \int_0^x (x-s)^{\alpha-1} z(s) ds \tag{5}$$

**Definition 4.** (Ajileye, James, Ayinde, & Oyedepo, 2022) Let  $y(x)$  be a continuous function, then

$${}_0I_x^\beta(y^{(\beta)}(x)) = y(x) - \sum_{k=0}^N \frac{y^{(k)}(0)}{k!} x^k \tag{6}$$

**Definition 5.** (Berinde, 2007) (Strict contraction) Let  $(X, d)$  be a metric space. A mapping  $T: X \rightarrow X$  is a strict contraction if  $T$  is  $\alpha$ -Lipschitzian with  $\alpha \in [0, 1)$ .

$$d(T(x), T(y)) \leq \alpha d(x, y) \ \forall x, y \in X$$

**Theorem 1.** (Berinde, 2007) (Banach fixed point theorem) Let  $(X, d)$  be a complete metric space and  $T: X \rightarrow X$  is a contraction on  $X$ , then  $T$  has a unique fixed point  $x \in X$  such that  $T(x) = x$ , and moreover the Picard successive approximations iterative scheme converges, given by

$$x_n = Tx_{n-1}$$

### 3. Materials and Methods

In this section, we establish the uniqueness of the solution and implement a collocation approach for the numerical solution of V-FIDEs.

**Lemma 1.** (Integral form): Let  $y \in C((0, 1), \mathbb{R})$  be the solution to equation (1), then it is equivalent to

$$y(x) + \frac{1}{\Gamma(\alpha)} \int_0^x (x-s)^{\alpha-1} (q(x)y(x)) ds + \frac{1}{\Gamma(\alpha)} \int_0^x (x-s)^{\alpha-1} \left( \lambda_1 \int_0^x k_1(x,t) F_1(y(t)) dt + \lambda_2 \int_0^1 k_2(x,t) F_2(y(t)) dt \right) ds = B(x) \tag{7}$$

where

$$B(x) = \sum_{k=0}^N \frac{y^{(k)}(0)}{k!} x^k + \frac{1}{\Gamma(\alpha)} \int_0^x (x-s)^{\alpha-1} g(s) ds$$

**Proof.**

Applying to equation (1) the operator  ${}_0I_x^\alpha(\cdot)$  gives

$${}_0I_x^\alpha \left( y'(x) \right) + \frac{1}{\Gamma(\alpha)} \int_0^x (x-s)^{\alpha-1} (q(x)y(x)) ds + \\ {}_0I_x^\alpha \left( \lambda_1 \int_0^x k_1(x,t) F_1(y(t)) dt + \lambda_2 \int_0^1 k_2(x,t) F_2(y(t)) dt \right) = {}_0I_x^\alpha (g(x))$$

and using equation (5) gives

$$y(x) + \frac{1}{\Gamma(\alpha)} \int_0^x (x-s)^{\alpha-1} (q(x)y(x)) ds + \frac{1}{\Gamma(\alpha)} \int_0^x (x-s)^{\alpha-1} \left( \lambda_1 \int_0^x k_1(x,t) F_1(y(t)) dt + \right. \\ \left. \lambda_2 \int_0^1 k_2(x,t) F_2(y(t)) dt \right) ds = \sum_{k=0}^N \frac{y^{(k)}(0)}{k!} x^k + \frac{1}{\Gamma(\alpha)} \int_0^x (x-s)^{\alpha-1} g(s) ds \\ y(x) + \frac{1}{\Gamma(\alpha)} \int_0^x (x-s)^{\alpha-1} (q(x)y(x)) ds + \frac{1}{\Gamma(\alpha)} \int_0^x (x-s)^{\alpha-1} \left( \lambda_1 \int_0^x k_1(x,t) F_1(y(t)) dt + \right. \\ \left. \lambda_2 \int_0^1 k_2(x,t) F_2(y(t)) dt \right) ds = W(x) \\ y(x) = W(x) - \frac{1}{\Gamma(\alpha)} \int_0^x (x-s)^{\alpha-1} (q(x)y(x)) ds - \frac{1}{\Gamma(\alpha)} \int_0^x (x-s)^{\alpha-1} \left( \lambda_1 \int_0^x k_1(x,t) F_1(y(t)) dt + \right. \\ \left. \lambda_2 \int_0^1 k_2(x,t) F_2(y(t)) dt \right) ds \tag{8}$$

where

$$W(x) = \sum_{k=0}^N \frac{y^{(k)}(0)}{k!} x^k + \frac{1}{\Gamma(\alpha)} \int_0^x (x-s)^{\alpha-1} g(s) ds$$

**3.1 Uniqueness of the method**

In order to establish the method's uniqueness, we provide the following hypothesis:

H<sub>1</sub>: There exist constants  $L_1, L_2 > 0$ , such that for any  $y_1, y_2 \in C([0,1], \mathbb{R})$

$$|F_1(y_1) - F_1(y_2)| \leq L_1 |y_1 - y_2| \\ \text{and} \\ |F_2(y_1) - F_2(y_2)| \leq L_2 |y_1 - y_2|$$

H<sub>2</sub>: There exist two functions  $k_1^*$  and  $k_2^* \in C([0,1] \times [0,1], \mathbb{R})$  in the set of all positive functions such that

$$k_1^* = \max_{x \in [0,1]} |\lambda_1| \int_0^x |k(x,t)| dt < \infty \\ k_2^* = \max_{x \in [0,1]} |\lambda_2| \int_0^1 |k(x,t)| dt < \infty$$

H<sub>3</sub>: The function  $g \in \mathbb{R}$  is continuous such that

$$q^* = \max_{x \in [0,1]} |q(x)|$$

**Theorem 2.** Let  $T: X \rightarrow X$  be the mapping defined by equation (8). Then  $T$  is a strict contraction if

$$\left( \frac{q^* + k_1^* + k_2^*}{\Gamma(\alpha + 1)} \right) < 1$$

**Proof.**

Let  $y_1(x), y_2(x) \in X$ . Applying Banach fixed point theorem to equation (8)

$$(Ty_1)(x) = W(x) - \frac{1}{\Gamma(\alpha)} \int_0^x (x-s)^{\alpha-1} (q(x)y_1(x)) ds - \frac{1}{\Gamma(\alpha)} \int_0^x (x-s)^{\alpha-1} \left( \lambda_1 \int_0^x k_1(x,t) F_1(y_1(t)) dt + \right. \\ \left. \lambda_2 \int_0^1 k_2(x,t) F_2(y_1(t)) dt \right) ds \tag{9}$$

and

$$(Ty_2)(x) = W(x) - \frac{1}{\Gamma(\alpha)} \int_0^x (x-s)^{\alpha-1} (q(x)y_1(x)) ds - \frac{1}{\Gamma(\alpha)} \int_0^x (x-s)^{\alpha-1} \left( \lambda_1 \int_0^x k_1(x,t) F_1(y_2(t)) dt + \lambda_2 \int_0^1 k_2(x,t) F_2(y_2(t)) dt \right) ds \tag{10}$$

Subtracting equation (10) from equation (9) gives

$$(Ty_1)(x) - (Ty_2)(x) = \frac{1}{\Gamma(\alpha)} \int_0^x (x-s)^{\alpha-1} [q(x)(y_2(x) - y_1(x))] ds + \frac{1}{\Gamma(\alpha)} \int_0^x (x-s)^{\alpha-1} \left( \lambda_1 \int_0^x k_1(x,t) F_1(y_2(t) - y_1(t)) dt + \lambda_2 \int_0^1 k_2(x,t) F_2(y_2(t) - y_1(t)) dt \right) ds$$

Taking the absolute value of both sides gives

$$|(Ty_1)(x) - (Ty_2)(x)| = \frac{1}{\Gamma(\alpha)} \int_0^x (x-s)^{\alpha-1} |q(x)(y_2(x) - y_1(x))| ds + \frac{1}{\Gamma(\alpha)} \int_0^x (x-s)^{\alpha-1} \left( |\lambda_1| \int_0^x |k_1(x,t)| |F_1(y_2(t) - y_1(t))| dt + |\lambda_2| \int_0^1 |k_2(x,t)| |F_2(y_2(t) - y_1(t))| dt \right) ds$$

Taking the maximum of both sides and applying  $H_1 - H_3$  gives  $d(Ty_1, Ty_2) \leq \left( \frac{q^* + k_1^* + k_2^*}{\Gamma(\alpha+1)} \right) d(y_1, y_2)$

Therefore T is a strict contraction mapping.

**Theorem 3.** (Continuity) Let  $(X, d)$  be a metric space and  $T: X \rightarrow X$  be a mapping, and let  $y_n(x), y(x) \in X$  and the  $\lim_{x \in [0,1]} y_n(x) = y(x)$ . Then T is continuous if  $d(Ty_n, Ty) \rightarrow 0$  as  $n \rightarrow \infty$ .

**Proof.**

$$\begin{aligned} |(Ty_n)(x) - (Ty)(x)| &= \frac{1}{\Gamma(\alpha)} \int_0^x (x-s)^{\alpha-1} |q(x)(y(x) - y_n(x))| ds + \\ &\frac{1}{\Gamma(\alpha)} \int_0^x (x-s)^{\alpha-1} \left( |\lambda_1| \int_0^x |k_1(x,t)| |F_1(y(t) - y_n(t))| dt + |\lambda_2| \int_0^1 |k_2(x,t)| |F_2(y(t) - y_n(t))| dt \right) ds \\ |(Ty_n)(x) - (Ty)(x)| &\leq \frac{1}{\Gamma(\alpha)} \int_0^x (x-s)^{\alpha-1} \max_{x \in [0,1]} |q(x)| \max_{x \in [0,1]} |y(x) - y_n(x)| ds + \\ &\frac{1}{\Gamma(\alpha)} \int_0^x (x-s)^{\alpha-1} \left( |\lambda_1| \max_{x \in [0,1]} \int_0^x |k_1(x,t)| \max_{x \in [0,1]} |F_1(y(t) - y_n(t))| dt + |\lambda_2| \max_{x \in [0,1]} \int_0^1 |k_2(x,t)| \max_{x \in [0,1]} |F_2(y(t) - y_n(t))| dt \right) ds \\ d(Ty_1, Ty_2) &\leq \left( \frac{q^* + k_1^* + k_2^*}{\Gamma(\alpha + 1)} \right) d(y_1, y_2) \end{aligned}$$

Since  $d(y_n, y) \rightarrow 0$  as  $n \rightarrow \infty$ , then  $d(Ty_n, Ty) \rightarrow 0$  as  $n \rightarrow \infty$ , therefore T is continuous.

### 3.2 Method of solution

Let the solution of equation (1) and equation (2) be approximated by

$$y(x) = \sum_{m=0}^M a_m x^m = \phi(x) \mathbf{A} \tag{11}$$

where  $\phi(x) = [1 \ x \ x^2 \ \dots \ x^M]$ ,  $\mathbf{A} = [a_0 \ a_1 \ \dots \ a_M]^T$

Substituting equation (11) into equation (8) gives

$$\phi(x) \mathbf{A} = W(x) - \frac{1}{\Gamma(\alpha)} \int_0^x (x-s)^{\alpha-1} (q(x)\phi(x)\mathbf{A}) ds - \frac{1}{\Gamma(\alpha)} \int_0^x (x-s)^{\alpha-1} \left( \lambda_1 \int_0^x k_1(x,t) F_1(\phi(t)\mathbf{A}) dt + \lambda_2 \int_0^1 k_2(x,t) F_2(\phi(t)\mathbf{A}) dt \right) ds \tag{12}$$

Collocating at  $x_i$  in equation (12) gives

$$\phi(x_i) \mathbf{A} = W(x_i) - \frac{1}{\Gamma(\alpha)} \int_0^x (x-s)^{\alpha-1} (q(x)\phi(x_i)\mathbf{A}) ds - \frac{1}{\Gamma(\alpha)} \int_0^x (x-s)^{\alpha-1} \left( \lambda_1 \int_0^x k_1(x,t) F_1(\phi(t)\mathbf{A}) dt + \lambda_2 \int_0^1 k_2(x,t) F_2(\phi(t)\mathbf{A}) dt \right) ds \tag{13}$$

Factorizing the value of  $\mathbf{A}$  from equation (13) gives

$$\left[ \left( \phi(x_i) + \frac{1}{\Gamma(\alpha)} \int_0^x (x-s)^{\alpha-1} (q(x)\phi(x_i)\mathbf{A}) ds + \frac{1}{\Gamma(\alpha)} \int_0^x (x-s)^{\alpha-1} \left( \lambda_1 \int_0^x k_1(x,t) F_1(\phi(t)\mathbf{A}) dt + \lambda_2 \int_0^1 k_2(x,t) F_2(\phi(t)\mathbf{A}) dt \right) ds \right) \mathbf{A} = W(x_i) \right] \quad (14)$$

Equation (14) can be in the form

$$J(x_i)\mathbf{A} = W(x_i) \tag{15}$$

where

$$J(x_i) = \phi(x_i) + \frac{1}{\Gamma(\alpha)} \int_0^x (x-s)^{\alpha-1} (q(x)\phi(x_i)\mathbf{A}) ds + \frac{1}{\Gamma(\alpha)} \int_0^x (x-s)^{\alpha-1} \left( \lambda_1 \int_0^x k_1(x,t) F_1(\phi(t)\mathbf{A}) dt + \lambda_2 \int_0^1 k_2(x,t) F_2(\phi(t)\mathbf{A}) dt \right) ds$$

We solve the system of equations (15) for the unknown values and substitute the results into the approximate solution to obtain the numerical result.

### 3.3 Convergence of the method

**Theorem 4. (Convergence of method)** Let  $(X, d)$  be a metric space and  $T: X \rightarrow X$  be a continuous mapping and  $y_N(x), y_{N-1}(x) \in X$  be approximate solutions of equation (7). Let  $\Delta_N(x) = |y_N(x) - y_{N-1}(x)|$ . If  $\lim_{N \rightarrow \infty} (\Delta_N(x)) \rightarrow 0$ , then the method converges to an exact solution.

**Proof.**

Let  $y_1(x)$  and  $y_2(x)$  be approximated by  $y_N(x) = \sum_{n=0}^M a_n x^n = \phi(x)\mathbf{A}$  and  $y_{N-1}(x) = \sum_{n=0}^M a_n x^n = \phi(x)\mathbf{B}$ . Substituting the approximate solution into equation (8) gives

$$Ty_N(x) = W(x) - \frac{1}{\Gamma(\alpha)} \int_0^x (x-s)^{\alpha-1} (q(x)\phi(x)\mathbf{A}) ds - \frac{1}{\Gamma(\alpha)} \int_0^x (x-s)^{\alpha-1} \left( \lambda_1 \int_0^x k_1(x,t) F_1(\phi(x)\mathbf{A}) dt + \lambda_2 \int_0^1 k_2(x,t) F_2(\phi(x)\mathbf{A}) dt \right) ds$$

Similarly

$$Ty_{N-1}(x) = W(x) - \frac{1}{\Gamma(\alpha)} \int_0^x (x-s)^{\alpha-1} (q(x)\phi(x)\mathbf{B}) ds - \frac{1}{\Gamma(\alpha)} \int_0^x (x-s)^{\alpha-1} \left( \lambda_1 \int_0^x k_1(x,t) F_1(\phi(x)\mathbf{B}) dt + \lambda_2 \int_0^1 k_2(x,t) F_2(\phi(x)\mathbf{B}) dt \right) ds$$

$$|(Ty_N)(x) - (Ty_{N-1})(x)| = \frac{1}{\Gamma(\alpha)} \int_0^x (x-s)^{\alpha-1} |q(x)\phi(x)| |\mathbf{B} - \mathbf{A}| ds + \frac{1}{\Gamma(\alpha)} \int_0^x (x-s)^{\alpha-1} \left( \lambda_1 \int_0^x k_1(x,t) |F_1(\phi(x)| |\mathbf{B} - \mathbf{A}| dt + \lambda_2 \int_0^1 k_2(x,t) |F_2(\phi(x)| |\mathbf{B} - \mathbf{A}| dt \right) ds$$

Since  $x \in [0,1]$  and  $|\mathbf{B} - \mathbf{A}| \neq 0$ , hence  $\lim_{N \rightarrow \infty} \Delta_N(x) \rightarrow 0$ . Therefore, the method converges.

### 3.4 Numerical examples

In this section, we provide numerical illustrations to assess the applicability and accuracy of the method. Let the approximate and exact solutions be  $y_n(x)$  and  $y(x)$  respectively.  $Error_N = |y_n(x) - y(x)|$ .

**Example 1.** (Hou *et al.*, 2021) Consider the first order V-FIDE

$$y'(x) + y(x) + \frac{1}{2} \int_0^x x y^2(t) dt - \frac{1}{4} \int_0^1 t y^3(t) dt = 2x + x^2 + \frac{1}{10} x^6 - \frac{1}{32} \tag{16}$$

subject to initial condition

$$y(0) = 0$$

The exact solution is  $y(x) = x^2$

**Solution 1.**

The approximate solution of equation (16) at  $N=5$  gives

$$y^5 = -2.546574062734 \times 10^{-15} - 2.842170943040 \times 10^{-14} x + 1.000000000000 x^2 + 1.818989403546 \times 10^{-12} x^3 + 1.818989403546 \times 10^{-12} x^4$$

**Example 2.** (Hou *et al.*, 2021) Consider the first order V-FIDE

$$y'(x) + y(x) - 2 \int_0^x \sin(x) y^2(t) dt = \cos(x) + (1 - x)\sin(x) + \cos(x)\sin^2(x) \tag{17}$$

subject to initial condition

$$y(0) = 0$$

The exact solution is  $y(x) = \sin(x)$

**Solution 2.**

The approximate solution of equation (17) at N=6 gives

$$y_6 = -4.737052972000 \times 10^{-10} + 1.000001048642x - 0.1394239556e - 2x^2 - 0.155098519841x^3 - 0.36024075147e - 1x^4 + 0.54202200699e - 1x^5 - 0.10605020725e - 1x^6.$$

**Example 3.** (Hou *et al.*, 2021) Consider the first order V-FIDE

$$y'(x) + \int_0^x (y^2(t) - 2) dt = \frac{1}{5}x^5 \tag{18}$$

subject to initial condition

$$y(0) = 0$$

The exact solution is  $y(x) = x^2$

**Solution 3.**

The approximate solution of equation (18) at N=4 gives

$$y_4 = -1.110223025000 \times 10^{-16}x + 1.000000000000x^2 - 3.552713679000 \times 10^{-15}x^4.$$

**4. Results and Discussion**

This section discusses the numerical results in the solved examples using the derived numerical approach. The result obtained for Example 1, as shown in Table 1, is that the approximate solution at  $N = 5$  namely  $y_5 = -2.546574062734 \times 10^{-15} - 2.842170943040 \times 10^{-14}x + 1.000000000000x^2 + 1.818989403546 \times 10^{-12}x^3 + 1.818989403546 \times 10^{-12}x^4$ .

The numerical result converged to an exact solution, and this confirms that our method performed better than the method proposed by (Hou *et al.*, 2021).

Table 1. Exact and approximate solutions, and absolute error for Example 1

x	Exact	Our method <sub>N=5</sub>	Errors	(Hou <i>et al.</i> , 2021) Error <sub>12</sub>
0.2	0.040000000000	0.040000000000	0.00	9.3974e-5
0.4	0.160000000000	0.160000000000	0.00	1.9813e-4
0.6	0.360000000000	0.360000000000	0.00	2.8891e-4
0.8	0.640000000000	0.640000000000	0.00	4.0786e-4
1.0	1.000000000000	1.000000000000	0.00	5.1647e-4

In numerical Example 2, as shown in Table 2, the approximate solution at  $N = 6$  is  $y_6 = -4.737052972000 \times 10^{-10} + 1.000001048642x - 0.1394239556e - 2x^2 - 0.155098519841x^3 - 0.36024075147e - 1x^4 + 0.54202200699e - 1x^5 - 0.10605020725e - 1x^6$ . The numerical result is better than the result obtained by (Hou *et al.*, 2021) at  $N = 12$ .

Table 2. Exact and approximate solutions, and absolute error for Example 2

x	Exact	Our method <sub>N=6</sub>	Errors	(Hou <i>et al.</i> , 2021) Error <sub>12</sub>
0.2	0.198669330800	0.198662679000	6.651800e-6	8.53966e-5
0.4	0.389418342300	0.389440411500	2.20692e-5	1.79180e-4
0.6	0.564642473400	0.564648677600	4.062042e-5	2.5968e-4
0.8	0.717356090900	0.717323556600	2.5674657-5	3.71839e-4
1.0	0.841470984800	0.841481394400	9.6104096e-5	5.01073e-4

The approximate solution obtained in Example 3 at  $N = 4$  is  $y_4 = -1.110223025000 \times 10^{-16}x + 1.000000000000x^2 - 3.552713679000 \times 10^{-15}x^4$ . The numerical result converged to an exact solution, and this confirms that our method performed better than the method proposed by (Hou *et al.*, 2021), as shown in Table 3.

Table 3. Exact and approximate solutions, and absolute error for Example 3

x	Exact	Our method <sub>N=4</sub>	Error <sub>4</sub>	(Hou <i>et al.</i> , 2021) Error <sub>4</sub>
0.2	0.040000000000	0.040000000000	0.00	4.6875e-17
0.4	0.160000000000	0.160000000000	0.00	2.77556e-17
0.6	0.360000000000	0.360000000000	0.00	1.11022e-16
0.8	0.640000000000	0.640000000000	0.00	2.22045e-16
1.0	1.000000000000	1.000000000000	0.00	1.11022e-15

## 5. Conclusions

For the numerical solution of the V-FIDEs, the collocation approach was investigated in this paper. This method is simple to apply, reliable, and efficient. For all computations in this work, Maple 18 was employed.

## References

- Agbolade, A. O. & Anake, T. A. (2017). Solution of first order Volterra linear integro-differential equations by collocation method. *Journal of Applied Mathematics*, Article ID. 1510267. doi:10.1155/2017/15267.
- Ahmed, A. H., Kirtiwant, P. G., & Shakir, M. A. (2019). The approximate solutions of fractional integro-differential equations by using modified adomian decomposition method. *Khayyan Journal of Mathematics*, 5(1), 21-39.
- Ahmad, N., Ullah, A., Ullah, A., Ahmad, S., Shah, K. & Ahmad, I. (2021). On analysis of the fuzzy fractional order Volterra-Fredholm integro-differential equation. *Alexandria Engineering Journal*, 60(1), 1827-1838.
- Ajileye, G. & Aminu, F. A. (2022). A numerical method using collocation approach for the solution of Volterra-Fredholm integro-differential equations. *African Scientific Reports*, 1, 205–211.
- Ajileye, G. & Aminu, F. A. (2022). Approximate solution to first-order integro-differential equations using polynomial collocation approach. *Journal Applied and Computational Mathematics*, 11, 486.
- Ajileye, G. & Amoo, S. A. (2023). Numerical solution to Volterra integro-differential equations using collocation approximation. *Mathematics and Computational Sciences*, 4(1), 1-8.
- Ajileye, G., James, A. A., Ayinde, A. M. & Oyedepo, T. (2022). Collocation approach for the computational solution of Fredholm-Volterra Fractional order of integro-differential equations. *Journal of the Nigerian Society Physical Science*, 4, 834.
- Amin, R., Shah, K., Mlaiki, N. Yuzabis, S., Abdeljawad T. & Hussain, A. (2022). Existence and numerical analysis using haar wavelet for fourth-order multi-term fractional differential equations. *Computational and Applied Mathematics*, 41, 329. Retrieved from <https://doi.org/10.1007/s40314-022-02041-8>.
- Atabakan, Z. P., Nasab, A. K., Kiliçman, A. & Eshkuvatov, Z. K. (2013). Numerical solution of nonlinear Fredholm integro-differential equations using spectral homotopy analysis method. *Mathematical Problems in Engineering*, 9(7), 120-130. Retrieved from <http://dx.doi.org/10.1155/2013/674364>.
- Berinde, V. (2007). *Iterative approximation of fixed points*. Bala Mare, Romania: Editura Efemeride.
- Bhrawy, A. H., Tohidi, E. & Soleymani, F. (2012). A new Bernoulli matrix method for solving high order linear and nonlinear Fredholm integro-differential equations with piecewise interval. *Applied Mathematics Computing*, 219, 482-497.
- Darania, P. & Ebadian, A. (2007). A method for the numerical solution of the integro-differential equations. *Applied Mathematics and Computation*, 188, 657–668.
- Ercan, C. & Kharerah, T. (2013). Solving a class of Volterra integral system by the differential transform method. *International Journal Nonlinear Science*, 16, 87-91.
- Garba, B. H. & Bichi, S. L. (2021). A hybrid method for solution of linear Volterra integro-differential equations via finite difference and Simpsons' numerical methods. *Open Journal Mathematics Analysis*, 5(1), 69-75.
- Hou, J., Niu, J., Xu, M. & Ngolo, W. (2021). A new numerical method to solve nonlinear Volterra-Fredholm integro-differential equations. *Mathematical Modelling and Analysis*, 26(3), 469-478.
- Irfan, N., Kumar, S. & Kapoor, S. (2014). Bernstein operational matrix approach for integro-differential equation arising in control theory. *Nonlinear Engineering*, 3, 117-123.
- Issa, K. & Saleh, F. (2017). Approximate solution of perturbed Volterra Fredholm integro differential equation by Chebyshev-galerkin method. *Journal of Mathematics*. doi:10.1155/2017/8213932.
- Khan, R. H. & Bakodah, H. O. (2013). Adomian decomposition method and its modification for nonlinear Abel's integral equations. *Computers and Mathematics with Applications*, 7, 2349-2358.
- Maadadi, A. & Rahmoune, A. (2018). Numerical solution of nonlinear Fredholm integro-differential equations using Chebyshev polynomials. *International Journal of Advanced Scientific and Technical Research*, 8(4), 85- 91. Retrieved from <https://dx.doi.org/10.26808/rs.st.i8v4.09>.

- Mehdiyera, G., Imanova, M., & Ibrahim, V. (2015). Solving Volterra Integro differential equation by second derivative methods. *Applied Mathematics Information Science*, 9, 2521 - 2527.
- Mehdiyeva, G., Ibrahimov, V., & Imanova, M. (2019). *On the construction of the multistep methods to solving the initial-value problem for ODE and the Volterra integro-differential equations*. Oxford, England: IAPE.
- Mittal, R. C. & Nigam, R. (2008). Solution of fractional integro-differential equations by Adomian decomposition method. *International Journal of Applied Mathematics and Mechanics*, 2, 87-94.
- Nemati, S., Lima, P., & Ordokhani, Y. (2015). Numerical method for the mixed Volterra-Fredholm integral equations using hybrid Legendre function. *Conference Application of Mathematics*, 4, 184-192.
- Rahman, M. (2007). *Integral equation and applications*. Boston, MA: WIT Press.
- Rahmani, L., Rahimi, B., & Mordad, M. (2011). Numerical solution of Volterra-Fredholm integro-differential equation by block pulse functions and operational matrices. *General Mathematics Notes*, 4(2), 37-48.
- Shahsavaran, A. & Shahsavaran, A. (2012). Application of Lagrange interpolation for nonlinear integro-differential equations. *Applied Mathematical Sciences*, 6(18), 887 - 892.
- Zada, L. Al-Hamami, M., Nawaz, R., Jehanzeb, S., Morsy, A., Abdel-Aty, A., & Nisar, K. S. (2021). A new approach for solving Fredholm integro-differential equations. *Information Sciences Getters*, 10(3), 3-10.