



Original Article

The χ^2 sequence space over p -metric spaces defined by Musielak modulus

Nagarajan Subramanian^{1*}, Chinappan Priya², and Nallamuthu Saivaraju²

¹ Department of Mathematics,
Shanmugha Arts, Science, Technology and Research Academy University, Thanjavur-613 401, India.

² Department of Mathematics,
Shri Angalamman College of Engineering and Technology, Trichirappalli-621 105, India

Received: 14 May 2014; Accepted: 1 August 2014

Abstract

In this paper, we introduce bonacci numbers of $\chi^2(F)$ sequence space over p -metric spaces defined by Musielak function and examine some topological properties of the resulting these spaces.

Keywords: analytic sequence, double sequences, χ^2 space, fibonacci number, Musielak - modulus function, p -metric space

1. Introduction

Throughout w , χ and Λ denote the classes of all, gai and analytic scalar valued single sequences, respectively.

We write w^2 for the set of all complex sequences (x_{mn}) , where $m, n \in \mathbb{N}$, the set of positive integers. Then, w^2 is a linear space under the coordinate-wise addition and scalar multiplication.

Some initial work on double sequence spaces is found in Bromwich(1965). Later on it was investigated by Hardy (1917); Moricz (1991); Moricz and Rhoades (1988); Basarir and Solankan (1999); Tripathy *et al.* (2003, 2004, 2006, 2007, 2008, 2009, 2010, 2011 and 2013); Turkmenoglu (1999); Raj (2010, 2011, 2012, 2013) and many others.

We procure the following sets of double sequences:

$$\mathcal{M}_u(t) := \left\{ (x_{mn}) \in w^2 : \sup_{m,n \in \mathbb{N}} |x_{mn}|^{t_{mn}} < \infty \right\},$$

$$\mathcal{C}_p(t) := \left\{ (x_{mn}) \in w^2 : p - \lim_{m,n \rightarrow \infty} |x_{mn} - l|^{t_{mn}} = 1 \text{ for some } l \in \mathbb{C} \right\},$$

$$\mathcal{C}_{0p}(t) := \left\{ (x_{mn}) \in w^2 : p - \lim_{m,n \rightarrow \infty} |x_{mn}|^{t_{mn}} = 1 \right\},$$

$$\mathcal{L}_u(t) := \left\{ (x_{mn}) \in w^2 : \sum_{m=1}^{\infty} \sum_{n=1}^{\infty} |x_{mn}|^{t_{mn}} < \infty \right\},$$

$$\mathcal{C}_{bp}(t) := \mathcal{C}_p(t) \cap \mathcal{M}_u(t) \text{ and } \mathcal{C}_{0bp}(t) = \mathcal{C}_{0p}(t) \cap \mathcal{M}_u(t);$$

where $t = (t_{mn})$ is the sequence of strictly positive reals t_{mn} for all $m, n \in \mathbb{N}$ and $p - \lim_{m,n \rightarrow \infty}$ denotes the limit in the Pringsheim's sense. In the case $t_{mn} = 1$ for all $m, n \in \mathbb{N}$; $\mathcal{M}_u(t), \mathcal{C}_p(t), \mathcal{C}_{0p}(t), \mathcal{L}_u(t), \mathcal{C}_{bp}(t)$ and $\mathcal{C}_{0bp}(t)$ reduce to the sets $\mathcal{M}_u, \mathcal{C}_p, \mathcal{C}_{0p}, \mathcal{L}_u, \mathcal{C}_{bp}$ and \mathcal{C}_{0bp} , respectively. Now, we may summarize the knowledge given in some document's related to the double sequence spaces. Gökhan *et al.* (2005) have proved that $\mathcal{M}_u(t)$ and $\mathcal{C}_p(t), \mathcal{C}_{bp}(t)$ are complete paranormed spaces of double sequences and obtained the α -, β -, γ - duals of the spaces $\mathcal{M}_u(t)$ and $\mathcal{C}_{bp}(t)$: Quite recently, in her PhD thesis, Zeltser (2001) has essentially studied both the theory of topological double sequence spaces and the theory of summability of double sequences. Mursaleen *et al.* (2003, 2004, 2013, 2014) and Tripathy *et al.* (2003, 2004, 2006, 2007, 2008, 2009, 2010, 2011 and 2013) have independently introduced the statistical convergence and Cauchy for double sequences and established the relation between statistical convergent and strongly Cesaro summable double sequences. Altay and Ba591sar (2005) have defined the spaces BS ; $BS(t)$, CS_p , CS_{bp} , CS_r and BV of

* Corresponding author.

Email address: nsmaths@yahoo.com

double sequences consisting of all double series whose sequence of partial sums are in the spaces M_u , $M_u(t)$, C_p , C_{bp} , C_r and L_u , respectively, and also examined some properties of those sequence spaces and determined the α - duals of the spaces BS , BV , CS_{bp} and the β (β) - duals of the spaces CS_{bp} and CS_r of double series. Ba592sar and Sever (2009) have introduced the Banach space L_q of double sequences corresponding to the well-known space ℓ_q of single sequences and examined some properties of the space \mathcal{L}_q . Recently Subramanian and Misra (2010) have studied the space $\chi_M^2(p, q, u)$ of double sequences and proved some inclusion relations.

The class of sequences which are strongly Cesaro summable with respect to a modulus was introduced by Maddox (1986) as an extension of the definition of strongly Cesaro summable sequences. Connor (1989) further extended this definition to a definition of strong A - summability with respect to a modulus where $A = (a_{n,k})$ is a nonnegative regular matrix and established some connections between strong A - summability, strong A - summability with respect to a modulus, and A - statistical convergence. In Pringsheim (1900) the four dimensional matrix transformation $(Ax)_{k,\ell} =$

$\sum_{m=1}^{\infty} \sum_{n=1}^{\infty} a_{k,m}^{mn} x_{mn}$ was studied extensively by Robison and Hamilton.

We need the following inequality in the sequel of the paper. For $a, b \geq 0$ and $0 < p < 1$; we have

$$(a+b)^p \leq a^p + b^p \quad (1.1)$$

The double series $\sum_{m,n=1}^{\infty} x_{mn}$ is called convergent if and only if the double sequence (s_{mn}) is convergent, where $s_{mn} = \sum_{i,j=1}^{m,n} x_{ij}$ ($m, n \in \mathbb{N}$).

A sequence $x = (x_{mn})$ is said to be double analytic if $\sup_{mn} |x_{mn}|^{1/m+n} < \infty$. The vector space of all double analytic sequences will be denoted by Λ^2 . A sequence $x = (x_{mn})$ is called double gai sequence if $((m+n)!|x_{mn}|)^{1/m+n} \rightarrow 0$ as $m, n \rightarrow \infty$. The double gai sequences will be denoted by χ^2 . Let $\phi = \{\text{all finite sequences}\}$.

Consider a double sequence $x = (x_{ij})$. The $(m, n)^{\text{th}}$ section $x^{[m,n]}$ of the sequence is defined by $x^{[m,n]} = \sum_{i,j=0}^{m,n} x_{i,j} \mathfrak{I}_{ij}$ for all $m, n \in \mathbb{N}$; where \mathfrak{I}_{ij} denotes the double sequence whose only non zero term is a $\frac{1}{(i+j)!}$ in the $(i, j)^{\text{th}}$ place for each $i, j \in \mathbb{N}$.

An FK-space(or a metric space) X is said to have AK property if (\mathfrak{I}_{mn}) is a Schauder basis for X . Or equivalently

$$x^{[m,n]} \rightarrow x.$$

An FDK-space is a double sequence space endowed with a complete metrizable; locally convex topology under which the coordinate mappings $x = (x_k) \rightarrow (x_{mn})$ ($m, n \in \mathbb{N}$) are also continuous.

Let M and Φ be mutually complementary modulus functions. Then, we have

(i) For all $u, y \geq 0$,

$$uy \leq M(u) + \Phi(y), \text{ (Young's inequality)}$$

[See Kampthan *et al.* (1981)] (1.2)

(ii) For all $u \geq 0$,

$$u\eta(u) = M(u) + \Phi(\eta(u)). \quad (1.3)$$

(iii) For all $u \geq 0$, and $0 < \lambda < 1$,

$$M(\lambda u) \leq \lambda M(u). \quad (1.4)$$

Lindenstrauss and Tzafriri (1971) used the idea of Orlicz function to construct Orlicz sequence space

$$\ell_M = \left\{ x \in w : \sum_{k=1}^{\infty} M\left(\frac{|x_k|}{\rho}\right) < \infty, \text{ for some } \rho > 0 \right\},$$

The space ℓ_M with the norm

$$\|x\| = \inf \left\{ \rho > 0 : \sum_{k=1}^{\infty} M\left(\frac{|x_k|}{\rho}\right) \leq 1 \right\},$$

becomes a Banach space which is called an Orlicz sequence space. For $M(t) = t^p$ ($1 \leq p < \infty$), the spaces ℓ_M coincide with the classical sequence space ℓ_p .

A sequence $f = (f_{mn})$ of modulus function is called a Musielak-modulus function. A sequence $g = (g_{mn})$ defined by

$$g_{mn}(v) = \sup \{ |v| u - f_{mn}(u) : u \geq 0 \}, m, n = 1, 2, \dots$$

is called the complementary function of a Musielak-modulus function f . For a given Musielak modulus function f , the Musielak-modulus sequence space t_f is defined by

$$t_f = \left\{ x \in w^2 : I_f(|x_{mn}|)^{1/m+n} \rightarrow 0 \text{ as } m, n \rightarrow \infty \right\},$$

where t_f is a convex modular defined by

$$I_f(x) = \sum_{m=1}^{\infty} \sum_{n=1}^{\infty} f_{mn} (|x_{mn}|)^{1/m+n}, x = (x_{mn}) \in t_f.$$

We consider t_f equipped with the Luxemburg metric

$$d(x, y) = \sup_{mn} \left\{ \inf \left(\sum_{m=1}^{\infty} \sum_{n=1}^{\infty} f_{mn} \left(\frac{|x_{mn}|^{1/m+n}}{mn} \right) \right) \leq 1 \right\}.$$

The notion of difference sequence spaces (for single sequences) was introduced by Kizmaz [16] as follows

$$Z(\Delta) = \{x = (x_k) \in w : (\Delta x_k) \in Z\},$$

for $Z = c, c_0$ and ℓ_∞ , where $\Delta x_k = x_k - x_{k+1}$ for all $k \in \mathbb{N}$

Here c, c_0 and ℓ_∞ denote the classes of convergent, null and bounded scalar valued single sequences respectively. The difference sequence space bv_p of the classical space ℓ_p is introduced and studied in the case $1 \leq p \leq \infty$ by Basar and Altay and in the case $0 < p < 1$ by Altay et al. (2005). The spaces $c(\Delta), c_0(\Delta), \ell_\infty(\Delta)$ and bv_p are Banach spaces normed by

$$\|x\| = |x_1| + \sup_{k \geq 1} |\Delta x_k| \text{ and } \|x\|_{bv_p} = \left(\sum_{k=1}^{\infty} |x_k|^p \right)^{1/p}, (1 \leq p < \infty).$$

Later on the notion was further investigated by many others. We now introduce the following difference double sequence spaces defined by

$$Z(\Delta) = \{x = (x_{mn}) \in w^2 : (\Delta x_{mn}) \in Z\},$$

where $Z = \Lambda^2, \chi^2$ and $\Delta x_{mn} = (x_{mn} - x_{mn+1}) - (x_{m+1n} - x_{m+1n+1}) = x_{mn} - x_{mn+1} - x_{m+1n} + x_{m+1n+1}$ for all $m, n \in \mathbb{N}$. The generalized difference double notion has the following representation: $\Delta^m x_{mn} = \Delta^{m-1} x_{mn} - \Delta^{m-1} x_{mn+1} - \Delta^{m-1} x_{m+1n} + \Delta^{m-1} x_{m+1n+1}$, and also this generalized difference double notion has the following binomial representation: $\Delta^m x_{mn} =$

$$\sum_{i=0}^m \sum_{j=0}^m (-1)^{i+j} \binom{m}{i} \binom{m}{j} x_{m+i, n+j}.$$

2. Definition and Preliminaries

Let $n \in \mathbb{N}$ and X be a real vector space of dimension w , where $n \leq w$. A real valued function $d_p(x_1, \dots, x_n) = \|(d_1(x_1, 0), \dots, d_n(x_n, 0))\|_p$ on X satisfying the following four conditions:

- (i) $\|(d_1(x_1, 0), \dots, d_n(x_n, 0))\|_p = 0$ if and only if $d_1(x_1, 0), \dots, d_n(x_n, 0)$ are linearly dependent,
- (ii) $\|(d_1(x_1, 0), \dots, d_n(x_n, 0))\|_p$ is invariant under permutation,
- (iii) $\|(\alpha d_1(x_1, 0), \dots, d_n(x_n, 0))\|_p = |\alpha| \|(d_1(x_1, 0), \dots, d_n(x_n, 0))\|_p, \alpha \in \mathbb{R}$
- (iv) $d_p((x_1, y_1), (x_2, y_2), \dots, (x_n, y_n)) = \left(d_X(x_1, x_2, \dots, x_n)^p + d_Y(y_1, y_2, \dots, y_n)^p \right)^{1/p}$ for $1 \leq p < \infty$;
(or)
- (v) $d((x_1, y_1), (x_2, y_2), \dots, (x_n, y_n)) = \sup \{d_X(x_1, x_2, \dots, x_n), d_Y(y_1, y_2, \dots, y_n)\}$, for $x_1, x_2, \dots, x_n \in X, y_1, y_2, \dots, y_n \in Y$ is called the p -product metric of the

Cartesian product of n -metric spaces is the p -norm of the n -vector of the norms of the n -sub spaces.

A trivial example of p -product metric of n -metric space is the p -norm space is $X = \mathbb{R}$ equipped with the following Euclidean metric in the product space is the p -norm:

$$\|(d_1(x_1, 0), \dots, d_n(x_n, 0))\|_p = \sup(|\det(d_{mn}(x_{mn}, 0))|) =$$

$$\sup \begin{pmatrix} d_{11}(x_{11}, 0) & d_{12}(x_{12}, 0) & \dots & d_{1n}(x_{1n}, 0) \\ d_{21}(x_{21}, 0) & d_{22}(x_{22}, 0) & \dots & d_{2n}(x_{1n}, 0) \\ \vdots & \vdots & \ddots & \vdots \\ d_{n1}(x_{n1}, 0) & d_{n2}(x_{n2}, 0) & \dots & d_{nn}(x_{nn}, 0) \end{pmatrix}$$

where $x_i = (x_{i1}, \dots, x_{in}) \in \mathbb{R}^n$ for each $i = 1, 2, \dots, n$.

If every Cauchy sequence in X converges to some $L \in X$, then X is said to be complete with respect to the p -metric. Any complete p -metric space is said to be p -Banach metric space.

2.1 Definition.

Let $A = (a_{k,\ell}^{mn})$ denote a four dimensional summability method that maps the complex double sequences x into the double sequence Ax where the k, ℓ -th term of Ax is as follows:

$$(Ax)_{k\ell} = \sum_{m=1}^{\infty} \sum_{n=1}^{\infty} a_{k\ell}^{mn} x_{mn}$$

such transformation is said to be non-negative if $a_{k\ell}^{mn}$ is non-negative.

The notion of regularity for two dimensional matrix transformations was presented by Silverman and Toeplitz. Following Silverman and Toeplitz, Robison and Hamilton presented the following four dimensional analog of regularity for double sequences in which both added an additional assumption of boundedness. This assumption was made since a double sequence which is P -convergent is not necessarily bounded.

Let λ and μ be two sequence spaces and $A = (a_{k,\ell}^{mn})$

be a four dimensional infinite matrix of real numbers $(a_{k,\ell}^{mn})$, where $m, n, k, \ell \in \mathbb{N}$. Then, we say A defines a matrix mapping from λ into μ and we denote it by writing $A : \lambda \rightarrow \mu$. if for every sequence $x = (x_{mn}) \in \lambda$ the sequence $Ax = \{(Ax)_{k\ell}\}$, the A -transform of x , is in μ . By $(\lambda : \mu)$, we denote the class of all matrices A such that $A : \lambda \rightarrow \mu$. Thus $A \in (\lambda : \mu)$ if and only if the series converges for each $k, \ell \in \mathbb{N}$. A

sequence x is said to be A - summable to α if Ax converges to α which is called as the A - limit of x .

2.2 Lemma. [See Maddox (1986)]

Matrix $A = (a_{k,\ell}^{mn})$ is regular if and only if the following three conditions hold:

(1) There exists $M > 0$ such that for every $k, \ell = 1, 2, \dots$

the following inequality holds: $\sum_{m=1}^{\infty} \sum_{n=1}^{\infty} |a_{k\ell}^{mn}| \leq M$;

(2) $\lim_{k,\ell \rightarrow \infty} a_{k\ell}^{mn} = 0$ for every $k, \ell = 1, 2, \dots$

(3) $\lim_{k,\ell \rightarrow \infty} \sum_{m=1}^{\infty} \sum_{n=1}^{\infty} a_{k\ell}^{mn} = 1$.

Let (q_{mn}) be a sequence of positive numbers and

$$Q_{k\ell} = \sum_{m=0}^k \sum_{n=0}^{\ell} q_{mn} \quad (k, \ell \in \mathbb{N}). \quad (2.1)$$

Then, the matrix $R^q = (r_{k\ell}^{mn})^q$ of the Riesz mean is given by

$$(r_{k\ell}^{mn})^q = \begin{cases} \frac{q_{mn}}{Q_{k\ell}} & \text{if } 0 \leq m, n \leq k, \ell \\ 0 & \text{if } (m, n) > k\ell \end{cases} \quad (2.2)$$

The fibonacci numbers are the sequence of numbers

$f_{k\ell}^{mn} (k, \ell, m, n \in \mathbb{N})$ defined by the linear recurrence equations $f_{00} = 1$ and $f_{11} = 1$, $f_{mn} = f_{m-1n-1} + f_{m-2n-2}$; $m, n \geq 2$. Fibonacci numbers have many interesting properties and applications in arts, sciences and architecture. Also, some basic properties of Fibonacci numbers are the following.

$$\sum_{k=1}^m \sum_{\ell=1}^n f_{mn} = f_{m+2n+2} - 1; m, n \geq 1,$$

$$\sum_{k=1}^m \sum_{\ell=1}^n f_{mn}^2 = f_{mn} f_{m+1n+1}; m, n \geq 1,$$

$$\sum_{k=1}^{\infty} \sum_{\ell=1}^{\infty} \frac{1}{f_{k\ell}^{mn}} \text{ converges.}$$

In this paper, we define the fibonacci matrix $F = (f_{k\ell}^{mn})_{m,n=1}^{\infty}$, which differs from existing Fibonacci matrix by using Fibonacci numbers $f_{k\ell}$ and introduce some new sequence spaces χ^2 and Λ^2 . Now, we define the Fibonacci matrix $F = (f_{k\ell}^{mn})_{m,n=1}^{\infty}$, by

$$(f_{k\ell}^{mn}) = \begin{cases} \frac{f_{k\ell}}{f_{(k+2)(\ell+2)} - 1} & \text{if } 0 \leq k \leq m; 0 \leq \ell \leq n \\ 0 & \text{if } (m, n) > k\ell \end{cases}$$

$$\begin{pmatrix} 1 & 0 & 0 & 0 & 0 \dots \\ \frac{1}{2} & \frac{1}{2} & 0 & 0 & 0 \dots \\ \frac{1}{4} & \frac{1}{4} & \frac{2}{4} & 0 & 0 \dots \\ \frac{1}{7} & \frac{1}{7} & \frac{2}{7} & \frac{3}{7} & 0 \dots \\ \vdots & \vdots & \vdots & \vdots & \ddots \end{pmatrix}.$$

It is obvious that the four dimensional infinite matrix F is triangular matrix. Also it follows from lemma 2.2 that the method F is regular.

Let M be an Musielak modulus function. We introduce the following sequence spaces based on the four dimensions infinite matrix F :

$$\begin{aligned} & \left[\Lambda_M^{2F}, \left\| (d(x_1, 0), d(x_2, 0), \dots, d(x_{n-1}, 0)) \right\|_p \right] = F_{\eta}(x) \\ & = \sup_{k\ell} \left\{ \sum_{m=1}^{\infty} \sum_{n=1}^{\infty} \left[M \left(f_{k\ell}^{mn} |x_{mn}|^{1/m+n}, \left\| (d(x_1, 0), d(x_2, 0), \dots, d(x_{n-1}, 0)) \right\|_p \right) \right] < \infty \right\} \\ & = \sup_{k\ell} \left\{ \frac{1}{f_{(k+2)(\ell+2)} - 1} \sum_{m=1}^{\infty} \sum_{n=1}^{\infty} \left[M \left(f_{k\ell}^{mn} |x_{mn}|^{1/m+n}, \left\| (d(x_1, 0), d(x_2, 0), \dots, d(x_{n-1}, 0)) \right\|_p \right) \right] < \infty \right\}, \\ & (k, \ell \in \mathbb{N}). \end{aligned}$$

Consider the metric space

$$\begin{aligned} & \left[\chi_M^{2F}, \left\| (d(x_1, 0), d(x_2, 0), \dots, d(x_{n-1}, 0)) \right\|_p \right] \text{ with the metric} \\ & d(x, y) = \sup_{k\ell} \left\{ M \left(F_{\eta}(x) - F_{\eta}(y) \right) : m, n = 1, 2, 3, \dots \right\}. \quad (2.3) \end{aligned}$$

$$\begin{aligned} & \left[\chi_M^{2F}, \left\| (d(x_1, 0), d(x_2, 0), \dots, d(x_{n-1}, 0)) \right\|_p \right] = F_{\mu}(x) \\ & = \lim_{k,\ell \rightarrow \infty} \\ & \left\{ \sum_{m=1}^{\infty} \sum_{n=1}^{\infty} \left[M \left(f_{k\ell}^{mn} ((m+n)! |x_{mn}|)^{1/m+n}, \left\| (d(x_1, 0), d(x_2, 0), \dots, d(x_{n-1}, 0)) \right\|_p \right) \right] = 0 \right\} \\ & = \lim_{m,n \rightarrow \infty} \\ & \left\{ \frac{1}{f_{(k+2)(\ell+2)} - 1} \sum_{m=1}^{\infty} \sum_{n=1}^{\infty} \left[M \left(f_{k\ell}^{mn} ((m+n)! |x_{mn}|)^{1/m+n}, \left\| (d(x_1, 0), d(x_2, 0), \dots, d(x_{n-1}, 0)) \right\|_p \right) \right] \right\} \\ & , (k, \ell \in \mathbb{N}). \end{aligned}$$

Consider the metric space

$$\begin{aligned} & \left[\chi_M^{2F}, \left\| (d(x_1, 0), d(x_2, 0), \dots, d(x_{n-1}, 0)) \right\|_p \right] \text{ with the metric} \\ & d(x, y) = \sup_{k\ell} \left\{ M \left(F_{\mu}(x) - F_{\mu}(y) \right) : m, n = 1, 2, 3, \dots \right\}. \quad (2.4) \end{aligned}$$

3. Main Results

3.1 Theorem.

The spaces $\left[\Lambda_M^{2F}, \|(d(x_1, 0), d(x_2, 0), \dots, d(x_{n-1}, 0))\|_p \right]$ and $\left[\chi_M^{2F}, \|(d(x_1, 0), d(x_2, 0), \dots, d(x_{n-1}, 0))\|_p \right]$ are BK spaces with the metric (2.4) and (2.5)

Proof: By Theorem 4.3.12 of Wilansky (1984) and since the four dimensional infinite matrix F is triangular, we have the result.

3.2. Theorem.

The spaces $\left[\Lambda_M^{2F}, \|(d(x_1, 0), d(x_2, 0), \dots, d(x_{n-1}, 0))\|_p \right]$ and $\left[\chi_M^{2F}, \|(d(x_1, 0), d(x_2, 0), \dots, d(x_{n-1}, 0))\|_p \right]$ are isomorphic to the spaces $\left[\Lambda_M^2, \|(d(x_1, 0), d(x_2, 0), \dots, d(x_{n-1}, 0))\|_p \right]$ and $\left[\chi_M^2, \|(d(x_1, 0), d(x_2, 0), \dots, d(x_{n-1}, 0))\|_p \right]$, respectively (i.e.) $\left[\Lambda_M^{2F}, \|(d(x_1, 0), d(x_2, 0), \dots, d(x_{n-1}, 0))\|_p \right] \cong \left[\Lambda_M^2, \|(d(x_1, 0), d(x_2, 0), \dots, d(x_{n-1}, 0))\|_p \right]$ and $\left[\chi_M^{2F}, \|(d(x_1, 0), d(x_2, 0), \dots, d(x_{n-1}, 0))\|_p \right] \cong \left[\chi_M^2, \|(d(x_1, 0), d(x_2, 0), \dots, d(x_{n-1}, 0))\|_p \right]$

Proof: Let us consider the space of χ^2 , since the four dimensional infinite matrix F is triangular, it has a unique inverse, which is also triangular [see Tripathy *et al.* (2009)]. Therefore the linear operator

$L_F : \left[\chi_M^{2F}, \|(d(x_1, 0), d(x_2, 0), \dots, d(x_{n-1}, 0))\|_p \right] \rightarrow \left[\chi_M^2, \|(d(x_1, 0), d(x_2, 0), \dots, d(x_{n-1}, 0))\|_p \right]$, defined by $L_F(x) = F(x)$ for all $x \in \left[\chi_M^{2F}, \|(d(x_1, 0), d(x_2, 0), \dots, d(x_{n-1}, 0))\|_p \right] \cong \left[\chi_M^2, \|(d(x_1, 0), d(x_2, 0), \dots, d(x_{n-1}, 0))\|_p \right]$,

is bijective and is metric preserving by (2.5) in Theorem 3.1. Hence

$\left[\chi_M^{2F}, \|(d(x_1, 0), d(x_2, 0), \dots, d(x_{n-1}, 0))\|_p \right] \cong$

$$\left[\chi_M^2, \|(d(x_1, 0), d(x_2, 0), \dots, d(x_{n-1}, 0))\|_p \right].$$

Similarly the proof for the other space can be established.

3.3 Theorem.

The inclusion

$$\left[\chi_M^2, \|(d(x_1, 0), d(x_2, 0), \dots, d(x_{n-1}, 0))\|_p \right] \subset$$

$$\left[\chi_M^{2F}, \|(d(x_1, 0), d(x_2, 0), \dots, d(x_{n-1}, 0))\|_p \right] \text{ and}$$

$$\left[\Lambda_M^2, \|(d(x_1, 0), d(x_2, 0), \dots, d(x_{n-1}, 0))\|_p \right] \subset$$

$$\left[\Lambda_M^{2F}, \|(d(x_1, 0), d(x_2, 0), \dots, d(x_{n-1}, 0))\|_p \right] \text{ holds.}$$

Proof: As F is a regular four dimensional infinite matrix, so the inclusion

$$\left[\chi_M^2, \|(d(x_1, 0), d(x_2, 0), \dots, d(x_{n-1}, 0))\|_p \right] \subset$$

$$\left[\chi_M^{2F}, \|(d(x_1, 0), d(x_2, 0), \dots, d(x_{n-1}, 0))\|_p \right] \text{ is obvious.}$$

Now, let $x = (x_{mn}) \in \left[\Lambda_M^{2F}, \|(d(x_1, 0), d(x_2, 0), \dots, d(x_{n-1}, 0))\|_p \right]$.

Then there is a constant $M > 0$ such that $|x_{mn}|^{1/m+n} \leq M$ for all $m, n \in \mathbb{N}$. Thus for each $k, \ell \in \mathbb{N}$.

$$|F_\eta(x)| \leq$$

$$\left\{ \frac{1}{f_{(k+2)(\ell+2)} - 1} \sum_{m=1}^k \sum_{n=1}^\ell \left[M \left(f_{k\ell}^{mn} |x_{mn}|^{1/m+n}, \|(d(x_1, 0), d(x_2, 0), \dots, d(x_{n-1}, 0))\|_p \right) \right] \right\}$$

$$\leq \left\{ \frac{M}{f_{(k+2)(\ell+2)} - 1} \sum_{m=1}^k \sum_{n=1}^\ell \left[M \left(f_{k\ell}^{mn} |x_{mn}|^{1/m+n}, \|(d(x_1, 0), d(x_2, 0), \dots, d(x_{n-1}, 0))\|_p \right) \right] \right\}$$

$< \infty$ which shows that $FX \in \left[\chi_M^2, \|(d(x_1, 0), d(x_2, 0), \dots, d(x_{n-1}, 0))\|_p \right]$.

$$\text{Thus we conclude that } \left[\chi_M^2, \|(d(x_1, 0), d(x_2, 0), \dots, d(x_{n-1}, 0))\|_p \right]$$

$$\subset \left[\chi_M^{2F}, \|(d(x_1, 0), d(x_2, 0), \dots, d(x_{n-1}, 0))\|_p \right].$$

Example: Consider the sequence $x = (x_{mn}) =$

$$\begin{pmatrix} 1 & 0 & 0 & 0 & 0 & \dots \\ 0 & 0 & 0 & 0 & 0 & \dots \\ 0 & 0 & 0 & 0 & 0 & \dots \\ 0 & 0 & 0 & 0 & 0 & \dots \\ \vdots & \vdots & \vdots & \vdots & \vdots & \vdots \end{pmatrix} \begin{pmatrix} -1 & 0 & 0 & 0 & 0 & \dots \\ 0 & 0 & 0 & 0 & 0 & \dots \\ 0 & 0 & 0 & 0 & 0 & \dots \\ 0 & 0 & 0 & 0 & 0 & \dots \\ \vdots & \vdots & \vdots & \vdots & \vdots & \vdots \end{pmatrix} \begin{pmatrix} 0 & 0 & 0 & 0 & 0 & \dots \\ 0 & 0 & 0 & 0 & 0 & \dots \\ 0 & 0 & 0 & 0 & 0 & \dots \\ 0 & 0 & 0 & 0 & 0 & \dots \\ \vdots & \vdots & \vdots & \vdots & \vdots & \vdots \end{pmatrix} \begin{pmatrix} 0 & 0 & 0 & 0 & 0 & \dots \\ 0 & 0 & 0 & 0 & 0 & \dots \\ 0 & 0 & 0 & 0 & 0 & \dots \\ 0 & 0 & 0 & 0 & 0 & \dots \\ \vdots & \vdots & \vdots & \vdots & \vdots & \vdots \end{pmatrix} \dots$$

Then we have for every $k, \ell \in \mathbb{N}$, $F_\mu(x) = \frac{1}{f_{(k+2)(\ell+2)} - 1} \sum_{m=1}^k \sum_{n=1}^\ell$

$$\left[M \left(f_{k\ell}^{mn} ((m+n)! |x_{mn}|)^{1/m+n}, \|(d(x_1, 0), d(x_2, 0), \dots, d(x_{n-1}, 0))\|_p \right) \right] \neq 0.$$

This shows that $FX \in \chi^2$ but x is not in χ^2 . Thus the

sequence x is in $\left[\chi_M^{2F}, \left\| (d(x_1, 0), d(x_2, 0), \dots, d(x_{n-1}, 0)) \right\|_p \right]$.

Hence the inclusion $\left[\chi_M^2, \left\| (d(x_1, 0), d(x_2, 0), \dots, d(x_{n-1}, 0)) \right\|_p \right]$

$\subseteq \left[\chi_M^{2F}, \left\| (d(x_1, 0), d(x_2, 0), \dots, d(x_{n-1}, 0)) \right\|_p \right]$ is strictly holds.

3.4 Theorem.

The sequence $x = (x_{mn}) \notin$

$\left[\chi_M^{2F}, \left\| (d(x_1, 0), d(x_2, 0), \dots, d(x_{n-1}, 0)) \right\|_p \right]$ but in

$\left[\Lambda_M^{2F}, \left\| (d(x_1, 0), d(x_2, 0), \dots, d(x_{n-1}, 0)) \right\|_p \right]$

Proof: Consider the sequence $x = (x_{mn})$

$= \begin{pmatrix} 1 & 1 & 1 & 1 & 1 \dots \\ 1 & 1 & 1 & 1 & 1 \dots \\ 1 & 1 & 1 & 1 & 1 \dots \\ 1 & 1 & 1 & 1 & 1 \dots \\ \vdots & \vdots & \vdots & \vdots & \vdots \end{pmatrix}$, for all $k, \ell \in \mathbb{N}$. Then we have for every

$k, \ell \in \mathbb{N}, F_\mu(x) = \begin{pmatrix} 1 & 1 & 1 & 1 & 1 \dots \\ 1 & 1 & 1 & 1 & 1 \dots \\ 1 & 1 & 1 & 1 & 1 \dots \\ 1 & 1 & 1 & 1 & 1 \dots \\ \vdots & \vdots & \vdots & \vdots & \vdots \end{pmatrix}$. This shows that

$FX \notin \left[\chi_M^{2F}, \left\| (d(x_1, 0), d(x_2, 0), \dots, d(x_{n-1}, 0)) \right\|_p \right]$.

Again, consider the sequence $x = (x_{mn}) = \frac{(-1)^{mn} (f_{m+2n+2+m+ln+1}) - 1}{f_{mn}}$,

for all $k, \ell \in \mathbb{N}$. Then we have for every $k, \ell \in \mathbb{N}, F_\eta(x) =$

$\begin{pmatrix} (-1)^{mn} & (-1)^{mn} \dots \\ \vdots & \vdots \end{pmatrix}$. This shows that

$FX \in \left[\Lambda_M^{2F}, \left\| (d(x_1, 0), d(x_2, 0), \dots, d(x_{n-1}, 0)) \right\|_p \right]$.

Competing Interests

Author has declared that no competing interests exist.

Acknowledgement

The authors thank the referee for his careful reading of the manuscript and comments that improved the presentation of the paper.

References

Altay, J. and Basar, F. 2005. Some new spaces of double sequences, *Journal of Mathematical Analysis and Applications*. 309(1), 70-90.

Basar, F. and Sever, Y. 2009. The space L_p of double sequences, *Mathematical Journal of Okayama University*. 51, 149-157.

Basarir, M. and Solancan, O. 1999. On some double sequence spaces, *Journal of Indian Academy of Mathematics*. 21(2), 193-200.

Bromwich, T.J.I'A. 1965. *An introduction to the theory of infinite series* Macmillan and Co. Ltd., New York.

Chandra, P. and Tripathy, B.C. 2002. On generalized Kothe-Toeplitz duals of some sequence spaces, *Indian Journal of Pure and Applied Mathematics*. 33(8), 1301-1306.

Cannor, J. 1989. On strong matrix summability with respect to a modulus and statistical convergence, *Canadian Mathematical Bulletin*. 32(2), 194-198.

Gökhan, A. and Colak, R. 2004. The double sequence spaces $c_2^p(p)$ and $c_2^{PB}(p)$, *Applied Mathematics Computation*. 157(2), 491-501.

Gökhan, A. and C596 olak, R. 2005. Double sequence spaces ℓ_2^∞ , *ibid*. 160(1), 147-153.

Gupta, M. and Pradhan, S. 2008. On Certain Type of Modular Sequence space, *Turkish Journal of Mathematics*. 32, 293-303.

Goes, G. and Goes, S. 1970. Sequences of bounded variation and sequences of Fourier coefficients, *Mathematische Zeitschrift*. 118, 93-102.

Hardy, G.H. 1917. On the convergence of certain multiple series, *Mathematical Proceedings of the Cambridge Philosophical Society*. 19, 86-95.

Hamilton, H.J. 1936. Transformations of multiple sequences, *Duke Mathematical Journal*. 2, 29-60.

Hamilton, H.J. 1938. A Generalization of multiple sequences transformation, *Duke Mathematical Journal*. 4, 343-358.

Hamilton, H.J. 1939. Preservation of partial Limits in Multiple sequence transformations, *Duke Mathematical Journal*. 4, 293-297.

Krasnoselskii, M.A. and Rutickii, Y.B. 1961. *Convex functions and Orlicz spaces*, Gorningen, Netherlands.

Kamthan, P.K. and Gupta, M. 1981. *Sequence spaces and series*, Lecture notes, Pure and Applied Mathematics, 65 Marcel Dekker, Inc., New York, U.S.A.

Kizmaz, H. 1981. On certain sequence spaces, *Canadian Mathematical Bulletin*. 24(2), 169-176.

Lindenstrauss, J. and Tzafriri, L. 1971. On Orlicz sequence spaces, *Israel Journal of Mathematics*. 10, 379-390.

Maddox, I.J. 1986. Sequence spaces defined by a modulus, Mathematical Proceedings of the Cambridge Philosophical Society. 100(1), 161-166.

Moricz, F. 1991. Extentions of the spaces c and c_0 from single to double sequences, *Acta Mathematica Hungarica*. 57(1-2), 129-136.

Moricz, F. and Rhoades, B.E. 1988. Almost convergence of double sequences and strong regularity of summability matrices, Mathematical Proceedings of the Cambridge Philosophical Society. 104, 283-294.

Mursaleen, M. and Edely, O.H.H. 2003. Statistical convergence of double sequences, *Journal Mathematical Analysis and Applications*. 288(1), 223-231.

Mursaleen, M. 2004. Almost strongly regular matrices and a core theorem for double sequences, *Journal Mathematical Analysis and Applications*. 293(2), 523-531.

Mursaleen, M. and Edely, O.H.H. 2004. Almost convergence and a core theorem for double sequences, *Journal Mathematical Analysis and Applications*. 293(2), 532-540.

Mursaleen, M., Sharma, S.K. and Kilicman, A. 2013. Sequence spaces defined by Musielak-Orlicz function over n -normed space, *Abstract and Applied Analysis*, Article ID 364743, p.10.

Mursaleen, M., Alotaibi, A. and Sharma, S.K. 2014. New classes of generalized seminormed difference sequence spaces, *Abstract and Applied Analysis*, Article ID 461081, p.7.

Mursaleen, M. and Sharma, S.K. 2014. Entire sequence spaces defined by Musielak-Orlicz function on locally convex Hausdorff topological spaces *Iranian Journal of Science and Technology. Transaction A*, 38.

Nakano, H., Concave modulars. 1953. *Journal of the Mathematical Society of Japan*. 5, 29-49.

Pringsheim, A. 1900. Zurtheorie der zweifach unendlichen zahlenfolgen, *Mathematische Annalen*. 53, 289-321.

Ruckle, W.H. 1973. FK spaces in which the sequence of coordinate vectors is bounded, *Canadian Journal of Mathematics*. 25, 973-978.

Robison, G.M. 1926. Divergent double sequences and series, *Transactions of the American Mathematical Society*. 28, 50-73.

Raj, K. and Sharma, S.K. 2011. Some sequence spaces in 2-normed spaces defined by Musielak Orlicz function, *Acta Universitatis Sapientiae Mathematica*. 3(1), 97-109.

Raj, K., Sharma, A.K. and Sharma, S.K. 2012. Sequence spaces defined by Musielak-Orlicz function in 2-normed spaces, *Journal of Computational Analysis and Applications*. Vol. 14.

Raj, K. and Sharma, S.K. 2013. Lacunary sequence spaces defined by Musielak-Orlicz function, *Le Matematiche*. 68, 33-51.

Raj, K., Sharma, A.K. and Sharma, S.K. 2011. A sequence spaces defined by a Musielak-Orlicz function, *International Journal of Pure and Applied Mathematics*. 67(4), 475-484.

Raj, K., Sharma, A.K. and Sharma, S.K. 2010. Difference sequence spaces in n -normed spaces defined by a Musielak-Orlicz function, *Armenian Journal of Mathematics*. 3, 127-141.

Raj, K. and Sharma, S.K. 2013. Ideal convergent sequence spaces defined by a Musielak-Orlicz function, *Thai Journal of Mathematics*. 11, 577-587.

Raj, K. and Sharma, S.K. 2012. Some new Lacunary strong convergent vector-valued multiplier difference sequence spaces defined by a Musielak-Orlicz function, *Acta Mathematica Academiae Nyiregyhaziensis*. 28, 103-120.

Subramanian, N. and Misra, U.K. 2010. The semi normed space defined by a double gai sequence of modulus function, *Fasciculi Mathematici*. 46.

Tripathy, B.C. and Mahanta, S. 2004. On a class of vector valued sequences associated with multiplier sequences, *Acta Math. Applicata Sinica (Eng. Ser.)*. 20(3), 487-494.

Tripathy, B.C. and Dutta, A.J. 2007. On fuzzy real-valued double sequence spaces ${}_2\ell_F^p$, *Mathematical and Computer Modelling*. 46 (9-10), 1294-1299.

Tripathy, B.C. and Sarma, B. 2008. Statistically convergent difference double sequence spaces, *Acta Mathematica Sinica*. 24(5), 737-742.

Tripathy, B.C. and Sarma, B. 2009. Vector valued double sequence spaces defined by Orlicz function, *Mathematica Slovaca*. 59(6), 767-776.

Tripathy, B.C. and Dutta, A.J. 2010. Bounded variation double sequence space of fuzzy real numbers, *Computers and Mathematics with Applications*. 59(2), 1031-1037.

Tripathy, B.C. and Sarma, B. 2011. Double sequence spaces of fuzzy numbers defined by Orlicz function, *Acta Mathematica Scientia*. 31 B(1)(2011), 134-140.

Tripathy, B.C. and Chandra, P. 2011. On some generalized difference paranormed sequence spaces associated with multiplier sequences defined by modulus function, *Analysis in Theory and Applications*. 27(1), 21-27.

Tripathy, B.C. and Dutta, A.J. 2013. Lacunary bounded variation sequence of fuzzy real numbers, *Journal in Intelligent and Fuzzy Systems*. 24(1), 185-189.

Tripathy, B.C. 2003. On statistically convergent double sequences, *Tamkang Journal of Mathematics*. 34(3), 231-237.

Tripathy, B.C. and Sen, M. 2006. Characterization of some matrix classes involving paranormed sequence spaces, *Tamkang Journal of Mathematics*. 37(2), 155-162.

Tripathy, B.C. and Hazarika, B. 2008. I -convergent sequence spaces associated with multiplier sequence spaces, *Mathematical Inequalities and Applications*. 11(3), 543-548.

Tripathy, B.C. and Chandra, P. 2011. On some generalized difference paranormed sequence spaces associated with multiplier sequences defined by modulus function, *Analysis in Theory and Applications*. 27(1), 21-27.

Tripathy, B.C. and Dutta, H. 2010. On some new paranormed difference sequence spaces defined by Orlicz functions, *Kyungpook Mathematical Journal*. 50(1), 59-69.

Turkmenoglu, A. 1999. Matrix transformation between some classes of double sequences, *Journal of Institute of Mathematics and Computer Science Maths Series*. 12(1), 23-31.

Woo, J.Y.T. 1973. On Modular Sequence spaces, *Studia Mathematica*. 48, 271-289.

Wilansky, A. 1984. *Summability through Functional Analysis*, North-Holland Mathematical Studies, North-Holland Publishing, Amsterdam. 85.

Zeltser, M. 2001., *Investigation of Double Sequence Spaces by Soft and Hard Analytical Methods*, *Dissertationes Mathematicae Universitatis Tartuensis* 25, Tartu University Press, University of Tartu, Faculty of Mathematics and Computer Science, Tartu.